

# Soil Microbial Responses to Lignitic Humic Acid Applications in Alkaline Soils

Dr. Safia Ali Khan, Dr. Muhammad Ali Shah, Dr. Rabia Nazir

Department of Environmental Sciences, University of Sindh, Jamshoro, Pakistan; Department of Soil and Environmental Sciences, Quaid-i-Azam University, Islamabad, Pakistan

## Abstract

**Indexed keywords:** Computer Engineering, Advanced Computing, Technology, Open Access

Article History: Received: 23 January 2022 | Accepted: 15 April 2022 | Published: 07 May 2022

A laboratory incubated study was conducted to evaluate the effects of lignitic humic acid addition on soil microbial activities and enzymatic reactions and indirectly improving the soil fertility level. Two types of soils (alkaline soils and sandy clay loam soils) were taken to apply lignitic humic acid at different rates (0, 2, 4, 6, 8  $\mu\text{g g}^{-1}$  soil). Treated and controlled soil samples were incubated at 25°C with moisture contents of 50% soil WHC for a period of 56 days. An increase in the cumulative CO<sub>2</sub>-C release, microbial biomass C, microbial biomass N, microbial biomass P and the activities of dehydrogenase and alkaline phosphatase enzymes in the soil was recorded by the addition of lignitic humic acid as compare to control. The decrease in ratios of C/N and C/P in the microbial biomass was observed, while the ratios of dehydrogenase/microbial biomass C, and alkaline phosphatase/microbial biomass C increased significantly. Addition of 8  $\mu\text{g g}^{-1}$  soil humic acid was the optimum humic acid application rate for its effect on soil microbial parameters in both soil types. All the microbial biomass indices exhibited similar temporal pattern being highest on 14<sup>th</sup> day of incubation followed by a gradual decline during rest of the incubation period. It was concluded that application of humic acid at an appropriate level (8  $\mu\text{g g}^{-1}$  soil) positively impacted soil ecology and soil fertility by improving microbial activities and enzymatic reactions in soil.

## Introduction

Deficiency of organic matter in soil has deleterious effects on soil properties and plant growth resulting in poor crop yields (Sarwar *et al.*, 2008; Hijbeek *et al.*, 2017). This challenging situation escalates the need to achieve the healthy soil status by using environmentally-friendly inputs for

sustaining crop production (Pukalchik *et al.*, 2019; Gazzola *et al.*, 2019). Humic substances form an important fraction of soil organic matter and are known to improve soil productivity by exerting favorable effects on soil properties such as aggregation, porosity, water retention, and cation exchange capacity (Hussein *et al.*, 2011; Muter *et al.*,



© 2022 The Authors. Open Access under CC BY 4.0.

How to cite: Dr. Safia Ali Khan, Dr. Muhammad Ali Shah, Dr. Rabia Nazir (2022). Soil Microbial Responses to Lignitic Humic Acid Applications in Alkaline Soils. Journal of Computer Engineering, 11(5), 77–93. DOI: <https://doi.org/10.5281/zenodo.19344818>

2015). Humic acid contribution to increase soil fertility on long term basis is highly recognized, as it helps in organic matter build up (Noroozisharaf and Kaviani, 2018), enhances uptake of macronutrients and micronutrients (Sharif *et al.*, 2002; Jones *et al.*, 2007; Eyheraguibel *et al.*, 2008), stimulates microbial growth, and promotes soil microbial activities (Kalaichelvi *et al.*, 2006, Valdrighi, 2013). Humic acid has a great tendency to involve in binding reactions which facilitates the substrate access to the active sites of the enzymes (Pukalchik *et al.*, 2019). Chen *et al.*, (2004) reported the chelating characteristics of humic acids to hold ionized nutrients to prevent the leaching losses of soil nutrients. Recently, the use of humic substances like sodium and potassium humates, are becoming more popular as an alternative to organic manures, to improve crop production in soils with low organic matter contents (Lee *et al.*, 2004; Tahir *et al.*, 2011, Arjumend *et al.*, 2015; Lyons and Genc, 2016; Kumar and Singh, 2017). Studies has shown that application of humic acid as an organic agricultural input can improve soil properties, nutrient availability and crop growth effectively, however, there is a lake of proper understanding of humic acid effects on microbial activities and the enzymatic reactions in soil, and rarely has been reported.

The activities of beneficial microbes in soil are vital for the long term sustainability of any type of soil and plant growth (García-Orenes *et al.*, 2013), as they play an integral role in the decomposition of organic matter (Wiseman *et al.*, 2012). Therefore, the status of soil microbial biomass content is considered as the most sensitive and rapid indicator of the long term soil productivity and a healthy soil ecosystem (Russell *et al.*, 2006; Smith *et al.*, 2007; Roger-Estrade *et al.*, 2010; García-Orenes *et al.*, 2013). Research has revealed that the soil microbial community can negatively be affected by soil

treatments by chemical applications (fertilizers/pesticides). Being the source and sink of essential plant nutrients, soil microbial biomass is a key element behind all biochemical transformations leading to soil nutrient availability (Nannipieri *et al.*, 2003) and can assist in measuring the status of processes like nitrogen fixation, nitrification, global carbon cycle (Wiseman *et al.*, 2012), solubilization of phosphate, and production of indole acetic acid (Bashan *et al.*, 2014; Viscardi *et al.*, 2016). Therefore, presence of beneficial soil microorganisms can possibly establish sustainable systems for productive soils (Coullerot *et al.*, 2013), and can be considered as biomarkers for soil fertility status by promoting mechanisms like biological fixation of nitrogen, solubilization of phosphate, production of indole acetic acid, etc. (Bashan *et al.*, 2014; Viscardi *et al.*, 2016). So far a few studies have been conducted to evaluate humic acid application effects on soil microbial activity (Filip *et al.*, 2004; Lizarazo *et al.*, 2005; Kalaichelvi *et al.*, 2006; Muscolo *et al.*, 2007), however, contradictory results are reported from most of these studies. Additionally, there are inconsistent references found for dose applications of humic acid to agricultural soils. The lack of understanding for humic acid effects on soil microbial biomass as well as its appropriate level of application to improve soil productivity demands a holistic consideration. It is very important to evaluate the effect of humic acid on microbial communities in the light of sustainable goals to improve or maintain the soil quality and biodiversity (García-Orenes *et al.*, 2013). The present study was conducted to investigate the effects of humic acid application at an optimum level on the size and activity of soil microbial biomass in relation to improve soil productivity on long term basis.

## Materials and methods

### Soils

Two alkaline soils were collected from two different locations in Pakistan with varying soil physical and chemical properties.

The sandy loam soil was collected from University Research Farm, Rawalpindi (referred as Rawalpindi soil), and the sandy clay loam soil was collected from University Research Farm, Koont (referred as Koont soil). The field moist soils were brought to the laboratory, hand-picked to remove stones, soil animals etc., passed through a 2-mm sieve, mixed thoroughly and frozen at  $-15^{\circ}\text{C}$  till the start of the experiment. A subsample of each soil was taken, air dried and mixed for the analysis of important physical and chemical soil properties, while the microbial parameters were measured in the field moist soil samples.

The Rawalpindi soil (sandy loam) was developed from sandstone while the Koont soil (sandy clay loam) was developed from the loess parent materials (Table 1). The Koont soil had higher sand, but lower contents of silt than the Rawalpindi soil, however both the soils had similar clay content. The Koont soil was also low in available P ( $< 3\mu\text{g P g}^{-1}$  soil), total N and organic C than the Rawalpindi soil, though both the soils had a similar pH and were alkaline in nature. All microbial parameters i.e., microbial biomass C, microbial biomass N, microbial biomass P, dehydrogenase and alkaline phosphatase activities were higher in the Rawalpindi soil as compared to the Koont soil.

**Table 1.** Physical and chemical properties of soils.

Property	Rawalpindi soil	Koont soil
Sand (%)	44.5	64.5
Silt (%)	32.5	15.0
Clay (%)	23.0	20.5
Texture	Loam	Sandy Clay Loam

pH <sub>1:2.5</sub>	7.9	8.2
EC 1:2.5 (dS m <sup>-1</sup> )	0.13	0.25
Organic C (mg g <sup>-1</sup> )	5.8	3.8
Total N (mg g <sup>-1</sup> )	0.5	0.3
Olsen P ( $\mu\text{g g}^{-1}$ )	3.6	2.0
Microbial biomass C ( $\mu\text{g g}^{-1}$ )	130.7	89.7
Microbial biomass N ( $\mu\text{g g}^{-1}$ )	11.3	6.9
Microbial biomass P ( $\mu\text{g g}^{-1}$ )	3.2	1.7
Dehydrogenase activity [TPF( $\mu\text{g g}^{-1}$ dwt 24 h <sup>-1</sup> )]	8.9	4.8
Alkaline phosphatase activity [ <i>p</i> -nitrophenol ( $\mu\text{g g}^{-1}$ dwt 24 h <sup>-1</sup> )]	45.7	20.1
C/N ratio	11.6	12.7

#### Lignitic humic acid

The lignitic humic acid used in the study was slightly acidic in nature (pH 6.0), had high contents of organic C (57.6%), and total N (4.2%).

The Olsen P contents in the humic acid were  $51.2\text{mg kg}^{-1}$ , while the extractable Fe, Cu, Mn and Zn contents were 6.5, 5.5, 14.1 and  $6.5\text{mg kg}^{-1}$ , respectively.

#### Experimental set up

The soil samples kept in the freezer were taken out, equilibrated to room temperature and the moisture contents were adjusted to 50% soil water holding capacity (WHC). The moist soil was then added to 2.0 l capacity incubation jars at the rate of  $600\text{g jar}^{-1}$  (on oven dry weight basis) and incubated at  $25^{\circ}\text{C}$  for 7 days (pre-incubation). After pre-incubation, the humic acid was applied at 0 (control),  $2\mu\text{g g}^{-1}$  soil,  $4\mu\text{g g}^{-1}$  soil,  $8\mu\text{g g}^{-1}$  soil and  $12\mu\text{g g}^{-1}$  soil and the jars were incubated for a period of 56 days at aforementioned conditions.  $\text{CO}_2\text{-C}$  evolution from each jar was measured after day 1, 2, 3, 5, 7, 10, 14 and thereafter weekly. The soil samples were collected from each incubation jar at day 0 (immediately after treatment application),

14, 28, 42 and 56 and analyzed for microbial biomass C, microbial biomass N, microbial biomass P, dehydrogenase activity and alkaline phosphatase activity.

#### Soil analysis

Soils were analyzed for important physical and chemical properties following the standard methods. Particle size analysis was carried out by Hydrometer method (Gee and Bauder, 1986). Water holding capacity (WHC) was determined by preparing saturated soil paste and measuring water contents gravimetrically in the saturated soil after surplus water removal (Anderson and Ingram, 1993).

Electrical conductivity of the soil saturation extract was recorded by conductivity meter (Rhoades, 1982) and the pH was measured in 1: 2.5 soil: water suspension with a standardized pH meter (McLean, 1982). Total organic C was determined by dichromate digestion method (Nelson and Sommers, 1982), total N was determined by Kjeldahl method (Bremner and Mulvaney, 1982), and Olsen P was determined in 0.5M NaHCO<sub>3</sub> (pH 8.5) soil extract using a spectrophotometer (Olsen and Sommers, 1982). For microbial analyses, the field moist soil samples stored at -15 °C were taken out of the freezer and equilibrated to room temperature.

Microbial biomass C and N were estimated by the fumigation-extraction method using 0.5M K<sub>2</sub>SO<sub>4</sub> as extractant (Brookes *et al.*, 1984; Vance *et al.*, 1987). Microbial biomass C was calculated as  $E_C / k_{EC}$ , where  $E_C$  = (organic C extracted from fumigated soils) - (organic C extracted from non-fumigated soils) and  $k_{EC}$  = 0.45 (Wu *et al.*, 1990). Microbial biomass N was calculated as  $E_N / k_{EN}$ , where  $E_N$  = (total N extracted from fumigated soils) - (total N extracted from nonfumigated soils) and  $k_{EN}$  = 0.54 (Brookes *et al.*, 1984). Soil microbial biomass P was also measured by the fumigation-extraction

method using 0.5M NaHCO<sub>3</sub> (pH 8.5) as extractant (Brookes *et al.*, 1982), as described by Joergensen *et al.* (1995). Microbial biomass P was calculated as  $E_P / k_{EP}$  / recovery, where  $E_P$  = (PO<sub>4</sub>-P extracted from fumigated soil) - (PO<sub>4</sub>-P extracted from non-fumigated soil) and  $k_{EP}$  = 0.40 (Brookes *et al.*, 1982). Dehydrogenase activity was measured by the reduction of 2,3,5-triphenyltetrazolium chloride in triphenylformazan (TPF) (Alef *et al.*, 1995) and alkaline phosphatase activity was measured using *p*-nitrophenyl phosphate as the substrate (Alef *et al.*, 1995; Nannipieri *et al.*, 2011). CO<sub>2</sub>-C evolved was trapped in 1.0 M NaOH, and measured by titration against standard HCl using phenolphthalein indicator (Anderson, 1982).

#### Analysis of Humic Acid

The pH and EC of the humic acid was measured in a 1:2 solid/water ratio (Peters *et al.*, 2003). Total organic C was determined by the Walkley-Black method (Nelson and Sommers, 1982). For the determination of total N and total P, the humic acid was digested in 1.2:1 H<sub>2</sub>SO<sub>4</sub>/H<sub>2</sub>O<sub>2</sub> mixture at 360 °C, and the total N and P in the digest were measured colorimetrically (Anderson and Ingram, 1993). Zinc, copper, iron and manganese contents in the digests were determined using atomic absorption spectrophotometer (GBC-932 plus).

#### Statistical analysis

The data obtained were analyzed statistically using Fisher's analysis of variance technique and the treatment means were compared by Tukey HSD Test at 5% level of significance (Steel *et al.*, 1997). Cumulative CO<sub>2</sub> release was analyzed by one-way ANOVA, while the microbial biomass C, microbial biomass N, microbial biomass P, dehydrogenase activity and alkaline phosphatase activity were analyzed by three-way ANOVA considering soils, humic acid levels and sampling days as the

factors. Software “Statstix 8.1” was employed for statistical analysis Russell and Eisensmith (1983).

## Results

### *Effect on soil microbial biomass*

Application of humic acid significantly promoted all the soil microbial biomass parameters as compared to control. The maximum increase was observed in the treatment receiving humic acid at  $8\mu\text{g g}^{-1}$  soil, followed by  $12\mu\text{g g}^{-1}$  soil humic acid treatment, which were significantly higher than other humic acid levels. However, Microbial biomass C, microbial biomass N and microbial biomass P contents varied significantly ( $p < 0.0001$ ) in the two soils (Table 2), being 1.5, 1.4 and 1.8 folds higher in the Rawalpindi soil than the Koont soil. The highest values of the above microbial parameters in both soils were measured at 14<sup>th</sup> day of incubation which declined significantly afterwards up to day 56 of incubation. This decline in microbial parameters was quite rapid during 14 to 28 days of the study period and slowed down later to become almost stable by the end of study (Fig. 1-6). Overall, the decline in microbial parameters was relatively more in Rawalpindi soil than the Koont soil.

**Table 2.** Properties of lignitic coal derived humic acid.

Property	Unit	Value
pH		6.1
Organic C	%	57.6
Total N	%	4.2
Olsen P	$\text{mg kg}^{-1}$	51.2
Zn	$\text{mg kg}^{-1}$	6.5
Mn	$\text{mg kg}^{-1}$	14.4
Fe	$\text{mg kg}^{-1}$	6.5
Cu	$\text{mg kg}^{-1}$	5.5

**Table 3.** Main effects of soils and humic acid application on microbial biomass indices of soils during 56 days of incubation.

The ratios of microbial biomass C/ N and microbial biomass C/ P were significantly higher in the Rawalpindi soil as compared to Koont soil.

Application of humic acid significantly reduced these ratios in both the soils. The maximum decline occurred with humic acid application at  $8\mu\text{g g}^{-1}$  soil, followed by  $12\mu\text{g g}^{-1}$  soil and  $4\mu\text{g g}^{-1}$  soil humic acid treatments which were statistically similar to each other. The lowest microbial biomass C/N and C/ P ratios were observed at day 14, which then increased gradually until the end of 56-day incubation.

### *Effect on enzymatic activities*

The results showed that humic acid treatments had significant effects on soil enzymatic reactions as compare to control. The activities of soil enzymes (dehydrogenase and alkaline phosphatase) were significantly ( $p < 0.0001$ ) different in the two soils

(Table 3). In the Rawalpindi soil, the dehydrogenase activity was 24% higher, and the alkaline phosphatase activity was 57% higher than in the Koont soil. Humic acid addition resulted in a significant 21% to 78% increase in dehydrogenase activity and 31% to 112% increase in alkaline phosphatase activity over the un-amended control. In both the cases, humic acid applied at  $8\mu\text{g g}^{-1}$  soil showed maximum increase in enzymes activities, followed by  $12\mu\text{g g}^{-1}$  soil humic acid and  $4\mu\text{g g}^{-1}$  soil humic acid application rates. The highest dehydrogenase activity was noticeably affected on the 0-day soil samples i.e., soil samples taken immediately after humic acid application, and showed a consistent decline thereafter throughout the incubation period. Contrary to dehydrogenase activity, the alkaline phosphatase activity increased during the first 14 days of incubation and thereafter declined till the end of incubation.

Main Effects	Microbial biomass C ( $\mu\text{g g}^{-1}$ soil)	Microbial biomass N ( $\mu\text{g g}^{-1}$ soil)	Microbial biomass P ( $\mu\text{g g}^{-1}$ soil)	Microbial biomass C/N	Microbial biomass C/P
Soils (S)					
Rawalpindi (S1)	151.2 A	13.6 A	4.12 A	11.3 A	44.9 A
Koont (S2)	102.4 B	9.4 B	2.308 B	11.1 B	37.6 B

	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001
LSD	0.723	0.075	0.038	0.073	0.39
D	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001
Humic acid levels (HA)					
S $\times$ HA	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001
S $\times$ D	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001
Control (T1)	111.8 D	8.9 E	2.5 E	12.7 A	46.1513 A
2 $\mu\text{g g}^{-1}$ soil (T2)	116.2 C	9.9 D	2.7 D	11.8 B	43.8309 B
4 $\mu\text{g g}^{-1}$ soil (T3)	128.1 B	11.06 C	3.390 C	10.7221 C	39.8300 C
8 $\mu\text{g g}^{-1}$ soil (T4)	139.6 A	13.8 A	3.9 A	10.1 D	37.2 D
12 $\mu\text{g g}^{-1}$ soil (T5)	138.1 A	12.9 B	3.6 B	C	39.5 C
LSD	1.603	0.166	0.083	0.163	0.876


## Sampling days (D)

Day 0	121.8 D	12.3 B	3.1 C	10.1 E	42.5 A
Day 14	135.9 A	12.7 A	3.7 A	10.1 D	38.7 C
Day 28	129.3 B	11.7 C	3.3 B	11.2 C	40.7 B
Day 42	124.9 C	10.6 D	3.1 C	11.8 B	41.5 B
Day 56	121.9 D	10.2 E	2.9 D	A	42.9 A
LSD	1.602	0.166	0.083	0.163	0.876


## Analysis of Variance

	<i>p</i> -value	<i>p</i> -value	<i>p</i> -value	<i>p</i> -value	<i>p</i> -value
S		<0.0001	<0.0001	<0.0001	<0.0001
HA Levels			<0.0001	<0.0001	<0.0001

\*Data are mean of three repeats. Means with different letters indicate significant difference at  $P < 0.05$  according to Tukey HSD Test.





**Fig. 1.** Microbial biomass C, N and P in (a) Rawalpindi and (b) Koont soils as affected by lignitic humic acid application during 56 days of incubation.




**Fig. 2.** Ratios of microbial biomass C/N and C/P in (a) Rawalpindi and (b) Koont soils as affected by lignitic humic acid application during 56 days of incubation.

**Fig. 1.** Microbial biomass C, N and P in (a) Rawalpindi and (b) Koont soils as affected by lignitic humic acid application during 56 days of incubation.



**Fig. 3.** Dehydrogenase and alkaline phosphatase activities in (a) Rawalpindi and (b) Koont soils as affected by lignitic humic acid application during 56 days of incubation.

The ratio of dehydrogenase/microbial biomass C was significantly lower in the Rawalpindi soil, while the ratio of alkaline phosphatase/microbial biomass C was significantly higher in the Rawalpindi soil as compared to Koont soil. Humic acid application significantly increased dehydrogenase/microbial biomass C (18.6% to 44%) and alkaline phosphatase/microbial biomass C (28% to 82%) ratios in both the soils, maximum at 8  $\mu\text{g g}^{-1}$  soil humic acid application rate. The maximum value of dehydrogenase/microbial biomass C was observed at day 0 of incubation, which decreased later throughout the incubation period (Fig.5). Unlike dehydrogenase/microbial biomass C, the ratio of alkaline phosphatase/microbial biomass C ratio increased significantly from 0 to 15 days of incubation, but thereafter declined till the end of incubation.



**Fig. 4.** Dehydrogenase/ microbial biomass C and alkaline phosphatase/microbial biomass C ratios in (a) Rawalpindi and (b) Koont soils as affected by lignitic humic acid application during 56 days of incubation.

*Effect on CO<sub>2</sub>-C evolution*

The mean CO<sub>2</sub>-C evolution rate from the soils, expressed as µg CO<sub>2</sub>-C g<sup>-1</sup> soil day<sup>-1</sup>, was significantly 2.6-fold higher in the Rawalpindi soil than the Koont soil (Table 4). Addition of humic acid increased mean CO<sub>2</sub>-C evolution from 16% to 73% in both the soils as compared to control, but maximum at humic acid application rate of 8 µg g<sup>-1</sup> soil. The cumulative CO<sub>2</sub>-C evolution during 56 days of incubation period was also significantly (1.9-fold) higher from the Rawalpindi soil as compared to that from the Koont soil (Table 5). There was 9% to 39% increase in cumulative CO<sub>2</sub>-C evolution from the soils at different levels of humic acid application, again being maximum at humic acid application rate of 8 µg g<sup>-1</sup> soil (Fig. 6).

**Table 4.** Main effects of soils and humic acid application on dehydrogenase (DHA), alkaline phosphatase (APA) activities and dehydrogenase/ microbial biomass C, and alkaline phosphatase/ microbial biomass C

ratios in the soils during 56 days of

	7.5	64.7			0.013
	0.371				
Sampling days (D)					
Day 0	9.3	A	45.8	E	0.08
Day 14	7.6	B	63.9	A	0.06
Day 28	5.9	C	59.3	B	0.05
Day 42	5.4	D	54.6	C	0.04
Day 56	4.7	E	49.6	D	0.04
LSD	0.371		1.675		0.002
Analysis of Variance					
	<i>p</i> -value	<i>p</i> -value	<i>p</i> -value	<i>p</i> -value	<i>p</i> -value
S	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001
HA	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001
D	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001
S × HA	0.0359	0.0005	<0.0001	<0.0001	<0.0001
S × D	<0.0001	0.9926	<0.0001	<0.0001	<0.0001
HA × D	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001
S × HA × D	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001
CV (±%)	7.88	4.28	6.19	4.43	

Data are mean of three repeats. Means with different

letters indicate significant difference at *P* < 0.05

according to Tukey HSD Test.

incubation.


DHA	APA	APA	
Main Effects [TPF (µg g <sup>-1</sup> 24 h <sup>-1</sup> )]			
	nitrophenol (µg g <sup>-1</sup> dwt h <sup>-1</sup> )]	DHA / Microbial C	Alkaline phosphatase / Microbial C

Soils (S)				
Rawalpindi (S1)	7.3	66.8	A	0.05
			B	0.44
			A	
Koont (S2)	5.9	42.5	B	0.06
			A	0.40
LSD	0.173	0.757	0.001	0.01
Humic acid levels (HA)				
Control (T1)	4.8	34.1	E	0.04
			D	0.29
2 µg g <sup>-1</sup> soil (T2)	5.8	44.6	D	0.05
			C	0.37
4 µg g <sup>-1</sup> soil (T3)	6.4	57.6	C	0.05
			C	0.45
8 µg g <sup>-1</sup> soil (T4)	8.5	72.3	A	0.06
			A	0.53
12 µg g <sup>-1</sup> soil (T5)	B		B	0.06
			B	0.47
LSD		1.675	0.002	

**Table 5.** Main effects of soils and humic acid application on rate and cumulative CO<sub>2</sub>-C evolution from soil during 56 days of incubation.

Main Effects	CO <sub>2</sub> -C evolution rate (μg CO <sub>2</sub> -C g <sup>-1</sup> soil day <sup>-1</sup> )	Σ evolution (μg CO <sub>2</sub> -C g <sup>-1</sup> soil)	CO <sub>2</sub> -C evolution (μg CO <sub>2</sub> -C g <sup>-1</sup> soil)
<b>Soils (S)</b>			
Rawalpindi	7.5	A	419.7
Koont	4.9	B	274.1
LSD	0.035		1.614
			6
<b>Humic acid levels (HA)</b>			
Control	4.4	E	207.5
2 μg g <sup>-1</sup> soil	5.1	D	226.5
4 μg g <sup>-1</sup> soil	6.3	C	253.5
8 μg g <sup>-1</sup> soil	7.6	A	287.9
12 μg g <sup>-1</sup> soil	7.5	B	280.8
LSD	0.0800		3.6725
<b>Analysis of Variance</b>			
	<i>p</i> -value		<i>p</i> -value
S	<0.0001		<0.0001
HA	<0.0001		<0.0001
S x HA	<0.0001		<0.0001
CV (±%)	0.74		0.84

Means with different letters indicate significant difference at  $P < 0.05$  according to Tukey HSD Test.



**Fig. 5.** CO<sub>2</sub>-C evolution from (a) Rawalpindi and (b) Koont soils as affected by lignitic humic acid application during 56 days of incubation.

## Discussion

This investigation concluded that addition of humic acid significantly improved the microbial biomass and enzymatic reactions. However, the response of the Rawalpindi soil to humic acid amendment was remarkably higher than koont soil in relation to all the soil microbial biomass parameters i.e., microbial biomass C, biomass N and biomass P. This attribute can of Rawalpindi soil can be because of higher contents of organic C, total N and available P in the Rawalpindi soil (Tajeda *et al.*, 2006; Malik *et al.*, 2013) as compare to Koont soil. Humic acid application significantly increased soil microbial biomass C. This indicates that humic acid stimulates the growth of microbial populations present in the soils (Muter *et al.*, 2015). The above observations are in line with the findings of Turgay *et al.* (2011) and Tavares & Nahas (2014), who reported increase in microbial biomass C by the addition of different organic materials, composts and humic substances to soils. However, no effects of humic acid on soil microbial biomass were observed by Little *et*

*al.* (2014). In the present study, microbial biomass C increased in the soils during first 14 days after humic acid application. Since both the soils had low organic matter contents, thus microbial populations in the soils were C restricted (Demoling *et al.*, 2007). Therefore, the increase in microbial biomass C can be attributed to the addition of a C source promoted the microbial populations in the soils. The decline in soil microbial biomass observed at later stages of incubation might be due to exhaustion of available carbon compounds with the time (Trevisan *et al.*, 2010; Malik *et al.*, 2013). Although humic acid is regarded as a highly decomposed and stable fraction of organic matter, yet it appears that the lignitic humic acid contains some organic components that can be utilized by the soil microorganisms as a source of energy.

Microbial biomass nitrogen is important because it controls soil organic nitrogen availability and loss, especially in high input systems (Moore *et al.*, 2000; Dong *et al.*, 2012). Humic acid application significantly increased microbial biomass N in both the soils. The maximum increase in microbial parameters occurred at  $8\mu\text{g g}^{-1}$  soil application rate, which seems to be an optimum level of humic acid application in the soils under study. These findings support the conclusion that there is always an optimum level for humic acid application (Sharif *et al.*, 2002; Haroon *et al.*, 2010). Although no reference exists for microbial biomass, but it has been shown that the excessive applications of humic acid can negatively affect plant growth (Atiyeh *et al.*, 2002), possibly through reduced availability of chelated nutrients (Chen *et al.*, 2004).

The assimilation of soil N by the increasing population of soil microorganisms occurred within first few hours of the humic acid application, and reached at its peak at 14th day of incubation. This immediate increase in microbial biomass N, as observed in case

of microbial biomass C as well, might be due to supplementation of C substrate as well as improving N availability to the soil microorganisms by humic acid (Tejada and Gonzalez, 2006). Microbial biomass N exhibited more temporal fluctuations than microbial biomass C. Similar temporal trend of microbial biomass N in comparison to microbial biomass C was reported by Joergensen (1995). In fact, soil microorganisms show great diversity in their N content than in their C content depending upon their stage of growth, and thus microbial biomass N always showed great variation due to diversity of soil microorganisms.

The increase in microbial biomass P might be ascribed to the growth of microbes which resulted in more assimilation of P into the microbial cells (Gichangi *et al.*, 2009). The temporal trend for variations in microbial biomass P was similar to that observed for microbial biomass C and microbial biomass N. In fact, the peaks of microbial biomass P at 14th day of incubation and its decline thereafter could be the result of microbial P mineralization as reported by others (Kabba *et al.*, 2004; Reddy *et al.*, 2005; Yang *et al.*, 2011; Yang *et al.*, 2013). Microbial biomass P is a highly labile source of P for plants because P assimilated in microbial cells is easily mineralized on microbial turn over (Reddy *et al.*, 2001; Reddy *et al.*, 2005).

Various studies are in agreement that humic acid generates an stimulatory effect on microorganisms (Kirschner *et al.*, 1999; Gryndler *et al.*, 2005; Tikhonov *et al.*, 2010; Kanaparthi and Conrad, 2015) by directly stimulation of biomass growth. This biomass growth occurs due to humic acid influence on biosynthetic activity that regulates the metabolism of a cell (Kirschner *et al.*, 1999; Kulikova *et al.*, 2005; Tikhonov *et al.*, 2010).

The net mineralization or immobilization of nitrogen in soils can be determined

primarily by carbon to nitrogen ratios of the added organic source. Soil microorganisms, particularly the bacteria and fungi, differ in their nitrogen requirements (Rashid *et al.*, 2016). Thus, microbial biomass C/ N ratio is frequently used to sense changes in the microbial community structure (Khan and Goergensen, 2009). The C/N ratio of microbial biomass in the two soils under study was almost similar. Humic acid application decreased the ratios of biomass C/ N in the soils indicating the role of humic acid in making N available to soil microorganisms. However, the decrease in C/ N ratio of microbial biomass might also be linked to the changes in microbial community structure caused by the humic acid application.

Addition of humic acid resulted in increase of dehydrogenase activity which is considered as a positive effect on soil health as soil enzymes are good markers of soil fertility being the biological catalysts in specific biochemical reactions. The activity of dehydrogenase has been proposed as a measure of the overall microbial activity in soils (Lizarazo *et al.*, 2005), and has also been indicated as a good index of soil microbial biomass in semi-arid Mediterranean areas (Garcia *et al.*, 1997). Therefore, humic acid application in relation to increased dehydrogenase activity in the soils can be attributed to its stimulation effect on the size and activity of soil microorganisms (Liang *et al.*, 2005; Haroon *et al.*, 2010). On an average, dehydrogenase activity was 34% higher in Rawalpindi soil than the Koont soil, which again be linked to the variation in organic C and microbial biomass contents in both the soils. The Rawalpindi soil had 2-folds higher organic C than the Koont soil, which resulted in high dehydrogenase activity in the former. It has been reported that incorporation of organic amendments to soil influences soil enzyme activities, because the added material may contain intra- and extra-cellular enzymes

and may also stimulate microbial activity in the soil (Pascual *et al.*, 2000).

Phosphatase is another important soil enzyme, because it provides P for plant uptake by the mineralization of soil organic P fractions. Being an extracellular hydrolase enzyme, it catalyzes the hydrolysis of phosphate from organic monoesters required by plants and microorganisms to maintain the cellular metabolism (Trevisan *et al.*, 2010; Malik *et al.*, 2013). Therefore, phosphatase is considered as an indicator of P mineralization potential of soils (Sinigani and Mahohi, 2009). In addition to dehydrogenase, the activity of alkaline phosphatase also increased in the soils due to humic acid application. The increasing demand for P by the growing microbial populations in the soils might have been responsible for the stimulation and the synthesis of phosphatase enzyme (Garcia *et al.*, 1996).

Addition of humic acid increased mean respiration rate as well as the cumulative CO<sub>2</sub>-C evolution from both the soils over the un-amended control.

These results are supported by the previous studies where humic acid application has been shown to immediately increase the soil respiration rate due to rapid promotion of the population of heterotrophic soil microorganisms. This happens due to the availability of carbon substrate following the organic amendments addition (Sarir *et al.*, 2006; Yang *et al.*, 2013). The fact that soil microbial biomass and soil respiration were promoted in lignitic humic acid amended soils indicates the ability of soil microbes to utilize some fraction of the organic compounds present in humic acid or in the soils (Cook and Allen, 1992). Additionally, the humic acid also contained essential elements like N, P and some micronutrients, therefore availability of these nutrients might have stimulated microbial populations in the soil (Piccolo, 2002).

Curves representing cumulative CO<sub>2</sub>-C with time showed that the slope at the outset was higher in the soil amended with humic acid suggesting that with the addition of this amendment, some fraction of the native soil organic carbon was also mineralized. Gilani and Behmanyar (2008) declared the availability of organic C as the most limiting factor for soil respiration. It is important to note that the CO<sub>2</sub> released from the soils may not merely come from the microbial respiration. The soils used in the present study were alkaline and had sufficient quantity of CaCO<sub>3</sub>, therefore, CO<sub>2</sub> released from action of humic acid on the soil carbonates may also contribute to the CO<sub>2</sub> release. However, the fact that the humic acid application had a similar effect on cumulative CO<sub>2</sub> release and on dehydrogenase activity, which is a measure of soil microbial activity, suggests that most of the CO<sub>2</sub> evolved did come from microbial respiration. It was observed that Soil respiration rates declined over time following cessation of labile carbon (C) inputs which can be attributed as a reduction of the available soil organic matter (SOM) pool as labile inputs diminish. This factor is in an agreement of several studies (Six & Jastrow, 2002; Conant *et al.*, 2011; Schmidt *et al.*, 2011).

### Conclusions

Application of humic acid significantly promoted the size and activity of soil microbial biomass, therefore acted as a soil biostimulant. However, the magnitude of increase in microbial biomass corresponded to the contents of native microbial biomass pool in the soils. Addition of 8µg g<sup>-1</sup> soil humic acid proved to be best recommended application rate to all other humic acid application levels. Temporal trend of soil respiration rate in response to humic acid application not only indicated the effect of humic acid on soil microbial CO<sub>2</sub> evolution but also specifies that most of the carbon utilized by microbes came from the native

organic C pool of the soils. Overall humic acid positive effect was recorded on microbial parameters including enzymatic reactions, and thus confirmed its stimulatory effect on microbial biomass. Therefore, humic acid application can be considered as an effective biotechnological tool for promotion of plant growth by developing a sustainable agriculture system.

### References

- Alef K, Nannipieri P, Trazar-Cepeda C.** 1995. Methods in applied soil microbiology and biochemistry. Academic, San Diego pp. 335-336.
- Anderson JM, Ingram JSI.** 1993. Tropical soil biology and fertility. CAB International, Wallingford
- Anderson JPE.** 1982. Soil respiration. Methods of soil analysis, part 2. Am Soc Agron, Soil Sci Soc Am, Madison Wisconsin pp. 831-871
- Arjumend T, Abbasi MK, Rafique E.** 2015. Effects of lignite-derived humic acid on some selected soil properties, growth and nutrient uptake of wheat (*Triticum aestivum* L.) grown under greenhouse conditions. Pak. J. Bot **47(6)**, 2231-2238
- Atiyeh RM, Edwards CA, Metzger JD, Lee S, Arancon NQ.** 2002. The influence of humic acids derived from earthworm-processed

- Balota EL, Collozi-Filho A, Andrade DS, Dick RP.** 2003. Microbial biomass in soils under different tillage and crop rotation systems. *Biology and Fertility of Soils* **38**, 15-20
- Bama KS, Selvakumari G, Santhi R, Singaram P.** 2003. Effect of humic acid on nutrient release pattern in an Alfisol (TypicHaplustalf). Tamil Nadu Agricultural University, Tamil Nadu, India. *Madras Agric. J* **90 (10-12)**, 665-670
- Banergee MR, Burton DL, McCaughey WP,**  
**Grant CA.** 2000. Influence of pasture management on soil biological quality. *Journal of Range Management* **53**, 127-133
- Bashan Y, de-Bashan LE, Prabhu SR, Hernandez JP.** 2014. Advances in plant growthpromoting bacterial inoculant technology: formulations and practical perspectives (1998-2013). *Plant Soil* **378**, 1-33.
- Bohn HL, McNeal BL, O'Connor GA.** 2001. *Soil Chemistry*. A Wiley Interscience Publ. John Wiley and Sons, New York
- Bremner JM, Mulvaney CS.** 1982. Methods of soil analysis, part 2 chemical and microbiological properties 595-624
- Brookes PC, Powlson DS, Jenkinson DS.** 1982. Measurement of microbial biomass phosphorus in soil. *Soil BiolBiochem* **14**, 319-329
- Brookes PC, Powlson DS, Jenkinson DS.** 1984.
- Phosphorus in the soil microbial biomass. *Soil Biol Biochem* **16**, 169-175
- Chen Y, De-Nobili M, Aviad T.** 2004. Stimulatory effect of humic substances on plant growth. Pp.103130. *Soil Organic Matter in Sustainable Agriculture*. CRC Press, Boca Raton, FL, USA
- Cimen F, Ok SS.** 2004. Properties of gyttja materials and their humic and fulvic acids in Afsin– Elbistan region. In *Proceedings of International Soil Congress on Natural Resource Management for Sustainable Development*, 7-10 June, Erzurum, Turkey pp. 34-38
- Conant RT, Ryanmg, Ågren GI, Birge HE, Davidson Ea, Eliasson PE, Evans SE, et al.** 2011. Temperature and soil organic matter decomposition rates - synthesis of current knowledge and a way forward. *Global Change Biology*, n/a–n/a.  
DOI:10.1111/j.1365-2486.2011.02496.x
- Couillerot O, Ramirez-Trujillo A, Walker V.** 2013. Comparison of prominent Azospirillum strains in Azospirillum-Pseudomonas-Glomus consortia for promotion of maize growth. *Appl. Microbiol. Biotechnol* **97(10)**, 4639-4649
- Demoling F, Figueroa D, Baath E.** 2007. Comparison of factors limiting bacterial growth in different soils. *Soil Biol-ogy and Biochemistry* **39**, 2485-2495
- Dong L, Córdova-Kreylos AL, Yang J, Yuan H, Scow KM.** 2012. Humic acids buffer the effects of urea on soil ammonia oxidizers and potential nitrification. *Soil BiolBiochem*



**Eyheraguibel B, Silvestre J, Morard P.** 2008. Effects of humic substances derived from organic waste enhancement on the growth and mineral nutrition of maize. *Bioresour. Technol* **99**, 4206-4212

**Filip Z, Tesarova M.** 2004. Microbial degradation and transformation of humic acid from permanent meadow and forest soils. *International Biodeterioration and Biodegradation* **54**, 225-231

**Garcia C, Hernandez T, Costa F, Barahona A.** 1996. Organic matter characteristics and nutrient content in eroded soils. *Environ. Manage* **20**, 133-141

**Garcia C, Hernandez T, Roldan A.** 1997. Changes in microbial activity after abandonment of cultivation in a semiarid Mediterranean environment. *J. Environ. Qual* **26**, 285-291

**García-Orenes F, Morugán-Coronado A, Zornoza R, Scow KM.** 2013. Changes in Soil Microbial Community Structure Influenced by Agricultural Management Practices in a Mediterranean Agro-Ecosystem **8(11)**, e80522.

DOI: 10.1371/journal.pone.0080522

**Gazzola P, Del Campo AG, Onyango A.** 2019. Going green vs going smart for sustainable development: Quo vadis? *J. Clean. Product* **214**, 881-

**892.**

**DOI:**

**10.1016/j.jclepro.2018.12.234**

**Gee GW, Bauder JW.** 1986. Particle size analysis. Pp.383-411. *Methods of Soil Analysis. Part-1.*, SSSA & ASA Monograph No. 9, Madison, Wisconsin, USA

**Gichangi EM, Mnkeni PNS, Brookes PC.** 2009.

Effects of goat manure and inorganic phosphate addition on soil inorganic and microbial biomass phosphorus fractions under laboratory incubation conditions

**Gilani SS, Bahmanyar MA.** 2008. Impact of Organic Amendments with and Without Mineral Fertilizers on Soil Microbial Respiration. *Journal of Applied Sciences* **8**, 642-647

**Gryndler M, Hršelová H, Sudová R, Gryndlerová H, Rezáčová V, Merhautová V.** 2005. Hyphal growth and mycorrhiza formation by the arbuscular mycorrhizal fungus *Glomus claroideum* BEG 23 is stimulated by humic substances. *Mycorrhiza* **15**, 483-488. DOI: 10.1007/s00572-005-0352-7 PubMed Abstract

| Cross Ref Full Text | Google Scholar

**Haroon R, Khattak RA, Muhammad D, Mian IA, Riaz M, Cresser MS.** 2010. Mechanism(s) of Humic Acid Induced Beneficial Effects in Saltaffected Soils. (Ph D Thesis). Department of Soil & Environmental Sciences, NWFP Agricultural University Peshawar, Pakistan

**Hijbeek R, Cormont A, Hazeu G, Bechini L, Zavattaro L, Janssen B, Werner M, Schlatter, N, Guzman G, Bijttebier J, Pronk AA, Eupen Mvan, Ittersum MKvan.** 2017. Do farmers perceive a deficiency of soil organic matter? A European and farm level analysis *Volume 83*,

**December 2017**, Pages 390-403

**Hussein K, Fawy HA.** 2011. Effect of Different Levels of Humic Acids on the Nutrient Content, Plant Growth, and Soil Properties under Conditions of Salinity. *Soil & Water Res* **6(1)**, 21-291 Soils and Water Department **Joergensen RG, Kubler H,**

- Meyer B, Wolters V.** 1995. Microbial biomass phosphorus in soils of beech (*Fagus sylvatica* L.) forests. *BiolFert Soils* **19**, 215-219
- Jones CA, Jacobsen JS, Mugaas A.** 2007. Effect of lowrate commercial humic acid on phosphorus availability, micronutrient uptake, and spring wheat yield. *Commun. Soil Sci. Plant Anal* **38**, 921-933
- Kabba BS, Aulakh MS.** 2004. Climatic conditions and crop-residue quality differentially affect N, P, and S mineralization in soils with contrasting P status. *J. Plant Nutr. Soil Sci* **167**, 596-601
- Kalaichelvi K, Chinnusamyl C, Swaminathan AA.** 2006. Exploiting the natural resource – lignite humic acid in agriculture - a review. *Agric. Rev* **27(4)**, 276 - 283
- Khan KS, Joergensen RG.** 2009. Changes in microbial biomass and P fractions in biogenic household waste compost amended with inorganic P fertilizers. *Bioresour Technol* **100**, 303-309
- Kirschner RA, Jr Parker BC, Falkinham JO.** 1999. Humic and fulvic acids stimulate the growth of *Mycobacterium avium*. *FEMS Microbiol. Ecol* **30**, 327-332. DOI: 10.1111/j.1574-6941.1999.tb00660.x  
PubMed Abstract | CrossRef Full Text | Google
- Kumar D, Singh AP.** 2017. Efficacy of Potassium Humate and Chemical Fertilizers on Yield and Nutrient Availability Patterns in Soil at Different Growth Stages of Rice, *Communications in Soil Science and Plant Analysis* **48(3)**, 245-261 DOI: 10.1080/00103624.2016.1261884.
- Lee JJ, Park RD, Kim YW, Shim, JH, Chae DH, Rim YS, Sohn BK, Kim TH, Kim KY.** 2004. Effect of food waste compost on microbial population, soil enzyme activity and lettuce growth. *Bioresource Technology* **93**, 21-28
- Liang Y, Nikolic M, Peng Y, Chen W, Jiang Y.** 2005. Organic manure stimulates biological activity and barley growth in soil subject to secondary salinization. *Soil Biol. Biochem* **37**, 1185-1195
- Little KR, Rose MT, Jackson WR, Cavagnaro T, Patti AF.** 2014. Do lignite-derived organic amendments improve early-stage pasture growth and key soil biological and physicochemical properties? *Crop and Pasture Science* **65**, 899-910
- Lizarazo LM, Jordá JD, Juárez M, SánchezAndreu J.** 2005. Effect of humic amendments on inorganic N, dehydrogenase, and alkaline phosphatase activities of a Mediterranean soil. *Biology and Fertility of Soils* **42**, 172-177
- Lyons G, Genc Y.** 2016. Commercial Humates in Agriculture: Real Substance or Smoke and Mirrors? *Agronomy* **6**, 1-8
- Mader P, Fließbach A, Dubois D, Gunst L, Fried P, Niggli U.** 2002. Soil fertility and biodiversity in organic farming. *Science* **296**, 1694-1697
- Malik MA, Khan KS, Marschner P, Ali S.** 2013. Organic amendments differ in their effect on microbial biomass and activity and on P pools in alkaline soils. *BiolFertl Soils* **49**, 415-425  
DOI: 10.1007/s00374-012-0738-6
- McLean EO.** 1982. Soil pH and lime requirement. *Methods of Soil Analysis, Part*

2: Chemical and Microbiological Properties. 2ed. Agronomy #9. pp. 199-224. American Society of Agronomy Inc, and Soil Science Society of America Inc. Madison, Wisconsin.

**Moore JM, Klose S, Tabatabai MA.** 2000. Soil microbial biomass carbon and nitrogen as affected by cropping systems. *Biology and Fertility of Soils* **31**, 200-210. DOI: 10.1007/s003740050646

**Muscolo A, Sidari M, Francioso O, Tugnoli V, Nardi S.** 2007. The auxin-like activity of humic substances is related to membrane interactions in carrot cell cultures. *J Chem Ecol* **33**, 115-129

**Muter O, Limane B, Strikauska S, Klavins M.** 2015. Effect of humic-rich peat extract on plant growth and microbial activity in contaminated soil. *Material Science and Applied Chemistry* **32**, 68-74

**Nannipieri P, Badalucco L.** 2003. Biological processes. *Handbook of Processes and Modelling in the Soil-Plant System*. Haworth Press, Binghamton, NY pp. 57-82

**Nannipieri P, Giagnoni L, Landi L, Renella G.** 2011. Role of phosphatase enzymes in soil. In: Bunemann EK, Oberson A, Frossard E (eds) *Phosphorus in action*. *Soil Biology* 26. Springer Verlag, Berlin pp 215-241

**Nelson DW, Sommers LE.** 1982. Total carbon, organic carbon and organic matter. *Methods of soil analysis. Part 2: chemical and microbiological properties*. American Society of Agronomy, Madison pp 539-579

**Noroozisharaf A, Kaviani M.** 2018. Effect of soil application of humic acid on nutrients uptake, essential oil and chemical compositions of garden thyme (*Thymus vulgaris* L.) under greenhouse conditions. *Physiol Mol Biol Plants* **24(3)**, 423-431.

Published online 2018 Mar 2. doi: 10.1007/s12298-

**Olsen SR, Sommers LE.** 1982. Phosphorus. p.

403-430. In A.L. Page (ed.) *Methods of soil analysis. Part 2. Chemical and microbiological properties*. 2nd ed. Agron.Monogr.no. 9. ASA and SSSA, Madison, WI

**Pascual JA, García C, Hernández T, Moreno JL, Ros M.** 2000. Soil microbial activity as a biomarker of degradation and remediation processes. *Soil Biology and Biochemistry* **32**, 1877-1883

**Peters J, Combs S, Hoskins B, Jarman J, Kovar J, Watson M, Wolf A, Wolf N.** 2003. *Recommended Methods of Manure Analysis (A3769)*; Cooperative Extension Publishing, University of Wisconsin-Extension: Madison, WI

**Piccolo A, Conte P, Trivellone E, Van Lagen B, Buurman P.** 2002. Reduced heterogeneity of a lignite humic acid by preparative HPSEC following interaction with an organic acid. Characterization of size-separates by PYRGC-MS and <sup>1</sup>H-NMR spectroscopy. *Environ. Sci. Technol* **36**, 76-84

**Pukalchik M, Kydralievva K, Yakimenko O, Fedoseeva E, Terekhova V.** 2019. Outlining the Potential Role of Humic Products in Modifying Biological Properties of the Soil—A Review. *Front. Environ. Sci.*, 07 June 2019. <https://doi.org/10.3389/fenvs.2019.00080>

**Rashid MI, Mujawar LH, Shahzad T, Almeelbi T, Ismail IMI, Oves M.** 2016. Bacteria and fungi can contribute to nutrients bioavailability and **Sarir MS, Durrani MI.** 2006. Utilization of natural resources for increase crop production. vol.

1, no. 2, August 2006 ISSN 1990-6145  
Journal of Agricultural and Biological  
Science ©2006 Asian Research  
Publishing Network (ARPN)

**Sarwar G, Schmeisky H, Hussain N,  
Muhammad S, Ibrahim M, Safdar S.**  
**183**, Pages 26-41

aggregate formation in degraded soils. Volume  
2008

**Reddy DD, Rao SA, Singh M.** 2001. Crop  
residue addition effects on myriad forms  
and sorption of phosphorus in a Vertisol.  
Bioresour Technol **80**, 93-99

**Reddy DD, Rao SA, Singh M.** 2005.  
Changes in P fractions and sorption in an  
Alfisol following crop residues application. J  
Plant Nutr Soil Sci **168**, 241-247

**Rhoades JD.** 1982. Soluble salts. p. 167–  
179. In A.L. Page (ed.) Methods of soil  
analysis. Part 2. Chemical and  
microbiological properties. 2nd ed. Chemical  
and microbiological  
properties. Agron. Monogr. no. 9. ASA and  
SSSA, Madison, WI.

**Robertson GP, Paul EA, Harwood RP.**  
2000. Greenhouse gases in intensive  
agriculture: Contributions of individual  
gases to the radiative forcing of the  
atmosphere. Science **289**, 1922-1925

**Roger-Estrade J, Anger C,  
Bertrand M,**

**Richard G.** 2010. Tillage and soil ecology:  
Partners for sustainable agriculture. Soils  
till Res **111**, 33-40.

DOI: 10.1016/j.still.2010.08.010

**Russel DF, Eisensmith SP.** 1983.  
MSTAT-C. Crop  
Soil Sci. Dept. Michigan State Univ. USA

2008. Improvement of Soil Physical and  
Chemical Properties with Compost  
Application in Rice-Wheat

**Russell L, Stokes AR, Macdonald H,  
Muscolo A, Nardi S.** 2006. Stomatal

Cropping System. Pak. J. Bot **40(1)**, 275-282,

responses to humic substances and auxin  
are sensitive to inhibitors of phospholipase  
A2. Plant Soil **283**, 175-185

**Schmidt MWI, Torn MS, Abiven S,  
Dittmar T, Guggenberger G, Janssens  
IA, Kleber M, et al.** 2011. Persistence of  
soil organic matter as an ecosystem property.  
Nature **478(7367)**, 49-56.  
DOI: 10.1038/nature10386

**Sharif M, Khattak RA, Sarir MS.** 2002.  
Effect of different levels of lignitic coal  
derived humic acid on growth of maize  
plants. Commun. Soil Sci. Plant Anal  
**33**, 3567-3580

**Sinegani AAS, Mahohi A.** 2009.  
Temporal variability of available P,  
microbial P and some  
phosphomonoesterase activities in a sewage  
sludge treated soil: The effect of soil water  
potential. African  
Journal of Biotechnology **8(24)**  
DOI: 10.4314/ajb.v8i24.68772

**Six J, Jastrow JD.** 2002. Organic matter  
turnover.  
Encyclopedia of soil science, 936-942.

**Smith JL, Il JM, Bolton Jr H, Bailey  
VL.** 2007. The initial rate of C substrate  
utilization and longer term soil C storage.  
Biol, Fert. Soils **44**, 315-320

**Steel RGD, Torrie JH, Dickey DA.** 1997. Principles and procedures of statistics; a biometrical approach. 3. Boston: McGraw-Hill;

**Sumayya SA, Siddique M, Mumtaz Ali K.** 2005. Study of the heavy metal pollution treatment potential of the coal generated humic acid. Journal of Basic and Applied Sciences **Vol 1**, No 2

**Tahir M, Khurshid MM, Khan MZ, Kazmi MH.**

2011. Lignite Derived Humic Acid Effect on Growth of Wheat Plants in Different Soils. Department of Soil and Environmental Sciences, Faculty of Agriculture, University of Azad Jammu and Kashmir, Rawalakot, Azad Jammu and Kashmir (Pakistan). Pedosphere

**Volume 21, Issue 1, February 2011**, Pages 124-131

**Tavares RLM, Nahas E.** 2014. Humic fractions of forest, pasture and maize crop soils resulting from microbial activity. Braz J Microbiol **45(3)**, 963-969.

**Tejada M, Gonzalez JL.** 2006. Effects of a crushed cotton gin compost on soil biological properties, nutrient leaching losses and maize yield. Agron J **98**, 749-759

**Tikhonov VV, Yakushev AV, Zavgorodnyaya YA, Byzov BA, Demin VV.** 2010. Effects of humic acids on the growth of bacteria. Eurasian Soil Sci **43**, 305-313. DOI:

10.1134/S1064229310030087

CrossRef Full Text | Google Scholar

**Trevisan S, Francioso O, Quaggiotti S, Nardi S.** 2010. Humic substances biological activity at the plant-soil interface Plant Signal Behav **5(6)**, 635-643

**Turgay OCA, Karaca S, Unver, Tamer N.** 2011. Effects of Coal- Derived Humic Substance on Some Soil Properties and Bread Wheat Yield, Communications in Soil Science and Plant Analysis

**42, 9**, 1050-1070

**Valdrighi MM, Pera A, Scatena S, Agnolucci M, Vallini G.** 2013. Effects of Humic Acids Extracted from Mined Lignite or Composted Vegetable Residues on Plant Growth and Soil Microbial Populations, Compost Science & Utilization **3**, 30-38

**Vance ED, Brookes PC, Jenkinson DS.** 1987. An extraction method for measuring soil microbial biomass C. Soil Biology and Biochemistry **19**, 703-

**707. DOI: 10.1016/0038-0717(87)90052-6**

**Viscardi S, Ventorino S, Duran P, Maggio A, De Pascale S, Mora ML, Pepe O.** 2016.

Assessment of plant growth promoting activities and abiotic stress tolerance of *Azotobacter chroococcum* strains for a potential use in sustainable agriculture. J. Soil Sci. Plant Nutr **16(3)**, 848-863

**Wiseman PE, Susan DD, Harris JR.** 2012. Organic Amendment Effects on Soil Carbon and Microbial Biomass in the Root Zone of Three Landscape Tree Species. Arboriculture & Urban Forestry 2012 **38(6)**, 262-276

**Wu J, Joergensen RG, Pommerening B,**

**Chaussod, R, Brookes PC.** 1990. Measurement of soil microbial biomass C by fumigation-extraction— an automated

procedure. *Soil Biology and Biochemistry*  
**Volume 22, Issue 8**, 1990, Pages  
1167-1169/. [https://doi.org/10.1016/0038-0717\(90\)](https://doi.org/10.1016/0038-0717(90)

**Yang KL, Guanc J, Zhuab, Yan L.** 2013  
Effects of Exogenous Humic Acids on Forms  
of Organic Phosphorus in Three Contrasting  
Types of Soil. *Communication s in Soilm  
Science and Plant Analysis* **44(14)**, 2095-  
2106, DOI: 10.1080/00103624.

**Yang LJ, Li TL, Li FS, Lemcoff JH.**  
2011. Longterm fertilization effect on  
fraction an