

Fixed Point Theorems for Generalized $(\alpha\beta-\psi)$ -Contractions in F-Metric Spaces: A Comprehensive Analysis

Ammar F. Al-Rashed, Muhammad Ahsan Raza, and Alessandro Bianchi

Department of Mathematical Sciences, King Fahad University of Petroleum and Minerals, Dhahran 31261, Saudi Arabia; (A.F.A.); (M.A.R.)

Abstract: The purpose of this paper is to define generalized $(\alpha\beta-\psi)$ -contraction in the context of F-metric space and obtain some new fixed point results. As applications, we solve a nonlinear neutral differential equation with an unbounded delay $\mathcal{G}'(t) = -\rho_1(t)\mathcal{G}(t) + \rho_2(t)L(\mathcal{G}(t - \zeta(t))) + \rho_3(t)\mathcal{G}'(t - \zeta(t))$, where $\rho_1(t), \rho_2(t)$ are continuous, $\rho_3(t)$ is continuously differentiable and $\zeta(t) > 0$, for all $t \in \mathbb{R}$ and is twice continuously differentiable.

Keywords: nonlinear neutral differential equation; fixed point; generalized $(\alpha\beta-\psi)$ -contraction; F-metric spaces

1. Introduction and Preliminaries

In 1906, M. Frechet introduced the notion of metric space [1], which is one pillar of not only mathematics but also physical sciences. Because of its importance and simplicity, this notion has been extended, improved and generalized in many different ways.

In 2018, Jleli et al. [2] introduced a fascinating generalization of metric space as follows:

Let $f \in \mathbf{F}$ and $f: (0, +\infty) \rightarrow \mathbb{R}$ be such that:

$$(F_1) \ 0 < \vartheta < \theta \implies f(\vartheta) \leq f(\theta),$$

$$(F_2) \ \text{for } \{\vartheta_n\} \subseteq \mathbb{R}^+, \lim_{n \rightarrow \infty} \vartheta_n = 0 \iff \lim_{n \rightarrow \infty} f(\vartheta_n) = -\infty.$$

Definition 1 ([2]). Let M be a nonempty set, and let $d_F: M \times M \rightarrow [0, +\infty)$ be a given mapping. Suppose that there exists $(f, h) \in \mathbf{F} \times [0, +\infty)$ such that

$$(D_1) \ (\vartheta, \theta) \in M \times M, \ d_F(\vartheta, \theta) = 0 \iff \vartheta = \theta.$$

$$(D_2) \ d_F(\vartheta, \theta) = d_F(\theta, \vartheta), \ \text{for all } (\vartheta, \theta) \in M \times M,$$

$$(D_3) \ \text{for every } (\vartheta, \theta) \in M \times M, \ N \in \mathbb{N}, \ N \geq 2, \ \text{and } (\vartheta_i)_{i=1}^N \subset \mathcal{M}, \ \text{with } (\vartheta_1, \vartheta_N) = (\vartheta, \theta), \ \text{we get}$$

$$N-1$$

$$d_F(\vartheta, \theta) > 0 \implies f(d_F(\vartheta, \theta)) \leq f\left(\sum_{i=1}^{N-1} d_F(\vartheta_i, \vartheta_{i+1})\right) + h.$$

Then d_F is said to be an F-metric on M , and the pair (M, d_F) is said to be an F-metric space.

Remark 1. They showed that any metric space is an F-metric space but the converse is not true in general, which confirms that this concept is more general than the standard metric concept.

Example 1 ([2]). The set of real numbers \mathbb{R} is an F-metric Space if we define d_F by

$$d_F(\vartheta, \theta) = \begin{cases} (\vartheta - \theta)^2 & \text{if } (\vartheta, \theta) \in [0, 3] \times [0, 3] \\ |\vartheta - \theta| & \text{if } (\vartheta, \theta) \notin [0, 3] \times [0, 3] \end{cases}$$

with $f(i) = \ln(i)$ and $h = \ln(3)$.

Definition 2 ([2]). Let (M, d_F) be an F-metric space.

(i) Let $\{\vartheta_n\}$ be a sequence in M . We say that $\{\vartheta_n\}$ is F-convergent to $\vartheta \in M$ if $\{\vartheta_n\}$ is convergent to ϑ with respect to the F-metric d_F . (ii) A sequence $\{\vartheta_n\}$ is F-Cauchy, if

$$\lim_{n, m \rightarrow \infty} d_F(\vartheta_n, \vartheta_m) = 0.$$

(iii) We say that (M, d_F) is F-complete, if every F-Cauchy sequence in M is F-convergent to a certain element in M .

Theorem 1 ([2]). Let (M, d_F) be an F-metric space and $H : M \rightarrow M$ be a given mapping. Suppose that the following conditions are satisfied:

- (i) (M, d_F) is F-complete, (ii) there exists $k \in (0, 1)$ such that

$$d_F(H(\vartheta), H(\theta)) \leq k d_F(\vartheta, \theta).$$

Then there exists $g^* \in M$ such that $Hg^* = g^*$ which is unique. Furthermore, for $\vartheta_0 \in M$, $\{\vartheta_n\} \subset M$ given by

$$\vartheta_{n+1} = H(\vartheta_n),$$

for all $n \in \mathbb{N}$, is F-convergent to g^* .

Afterwards, Hussain et al. [3] considered the notion of α - ψ -contraction in the setting of F-metric spaces and proved the following fixed point theorem.

Theorem 2 ([3]). Let (M, d_F) be an F-metric space and $H : M \rightarrow M$ be β -admissible mapping. Suppose that the following conditions are satisfied:

- (i) (M, d_F) is F-complete,
- (ii) there exists $\beta : M \times M \rightarrow [0, +\infty)$ and $\psi \in \Psi$ such that

$$\beta(\vartheta, \theta) d_F(H(\vartheta), H(\theta)) \leq \psi(M(\vartheta, \theta)),$$

where

$$M(\vartheta, \theta) = \max \{d_F(\vartheta, \theta), d_F(\vartheta, H\vartheta), d_F(\theta, H\theta)\},$$

for all $\vartheta, \theta \in \mathbf{M}$,

(iii) there exists $\vartheta_0 \in \mathbf{M}$ such that $\beta(\vartheta_0, H(\vartheta_0))$. Then there exists unique $\vartheta^* \in \mathbf{M}$ such that $H\vartheta^* = \vartheta^*$.

For more details in this direction, we refer the readers to References [4–10].

On the other hand, Samet et al. [11] introduced the concepts of α - ψ -contractive and α -admissible mappings and established various fixed point theorems for such mappings in complete metric spaces.

Denote with Ψ the family of nondecreasing functions $\psi: [0, +\infty) \rightarrow [0, +\infty)$ such that $\sum_{n=1}^{\infty} \psi^n(\vartheta) < +\infty$ for all $\vartheta > 0$, where ψ^n is the n -th iterate of ψ .

The following lemma is well known.

Lemma 1. *If $\psi \in \Psi$, then the following hold:*

(i) $(\psi^n(\vartheta))_{n \in \mathbf{N}}$ converges to 0 as $n \rightarrow \infty$ for all $\vartheta \in (0, +\infty)$,

(ii) $\psi(\vartheta) < \vartheta$ for all $\vartheta > 0$, (iii) $\psi(\vartheta) = 0$ iff $\vartheta = 0$.

Samet et al. [11] defined the notion of α -admissible mappings as follows:

Definition 3 ([11]). *Let H be a self-mapping on \mathbf{M} and $\alpha: \mathbf{M} \times \mathbf{M} \rightarrow [0, +\infty)$ be a function. We say that H is an α -admissible mapping if*

$$\alpha(\vartheta, \theta) \geq 1 \implies \alpha(H\vartheta, H\theta) \geq 1,$$

for all $\vartheta, \theta \in \mathbf{M}$.

Hussain et al. [12] extended the above notion of α -admissible mapping as follows.

Definition 4 ([12]). *Let H be a self-mapping on \mathbf{M} and $\alpha, \beta: \mathbf{M} \times \mathbf{M} \rightarrow [0, +\infty)$ be two functions. We say that H is an α -admissible mapping with respect to β if*

$$\alpha(\vartheta, \theta) \geq \beta(\vartheta, \theta) \implies \alpha(H\vartheta, H\theta) \geq \beta(H\vartheta, H\theta),$$

for all $\vartheta, \theta \in \mathbf{M}$.

If $\beta(\vartheta, \theta) = 1$, then Definition 4 reduces to Definition 3.

Later on, the authors (see References [13,14]) utilized the above concepts and obtained different fixed point results.

In this paper, we define the notion of generalized $(\alpha\beta\text{-}\psi)$ -contraction and establish some new fixed point theorems in the context of F -metric spaces. We also furnish a notable example to describe the significance of established results.

2. Results and Discussions

Definition 5. *Let (\mathbf{M}, d_F) be an F -metric space and $H: \mathbf{M} \rightarrow \mathbf{M}$. Then H is said to be generalized $(\alpha\beta\text{-}\psi)$ -contraction if there exists $\alpha, \beta: \mathbf{M} \times \mathbf{M} \rightarrow [0, \infty)$ and $\psi \in \Psi$ such that $\alpha(\vartheta, H\vartheta)\alpha(\theta, H\theta) \geq \beta(\vartheta, H\vartheta)\beta(\theta, H\theta)$ implies*

$$d_F(H\vartheta, H\theta) \leq \psi(\max\{d_F(\vartheta, \theta), \min\{d_F(\vartheta, H\vartheta), d_F(\theta, H\theta)\}\}), \quad (1)$$

for all $\vartheta, \theta \in \mathbf{M}$.

Theorem 3. Let (\mathbf{M}, d_F) be an F-metric space and let $H : \mathbf{M} \rightarrow \mathbf{M}$ be generalized $(\alpha\beta-\psi)$ -contraction. Suppose that the following assertions hold:

- (i) (\mathbf{M}, d_F) is F-complete,
- (ii) H is an α -admissible mapping with respect to β ,
- (iii) there exists $\vartheta_0 \in \mathbf{M}$ such that $\alpha(\vartheta_0, H\vartheta_0) \geq \beta(\vartheta_0, H\vartheta_0)$,
- (v) either H is continuous or if $\{\vartheta_n\}$ is a sequence in \mathbf{M} such that $\vartheta_n \rightarrow \vartheta$, $\alpha(\vartheta_n, \vartheta_{n+1}) \geq \beta(\vartheta_n, \vartheta_{n+1})$,

then $\alpha(\vartheta, H\vartheta) \geq \beta(\vartheta, H\vartheta)$.

Then there exists $\vartheta^* \in \mathbf{M}$ such that $\vartheta^* = H\vartheta^*$.

Proof. Let $\vartheta_0 \in \mathbf{M}$ be such that $\alpha(\vartheta_0, H\vartheta_0) \geq \beta(\vartheta_0, H\vartheta_0)$ and construct $\{\vartheta_n\}$ in \mathbf{M} by $\vartheta_{n+1} = H^n\vartheta_0 =$

$H\vartheta_n, \forall n \in \mathbf{N}$. By (ii), we have

$$\alpha(\vartheta_0, \vartheta_1) = \alpha(\vartheta_0, H\vartheta_0) \geq \beta(\vartheta_0, H\vartheta_0) = \beta(\vartheta_0, \vartheta_1).$$

Continuing in this way, we get

$$\alpha(\vartheta_{n-1}, \vartheta_n) = \alpha(\vartheta_{n-1}, H\vartheta_{n-1}) \geq \beta(\vartheta_{n-1}, H\vartheta_{n-1}) = \beta(\vartheta_{n-1}, \vartheta_n), \quad (2)$$

for all $n \in \mathbf{N}$. Then

$$\alpha(\vartheta_{n-1}, H\vartheta_{n-1})\alpha(\vartheta_n, H\vartheta_n) \geq \beta(\vartheta_{n-1}, H\vartheta_{n-1})\beta(\vartheta_{n-1}, H\vartheta_{n-1}), \quad (3)$$

for all $n \in \mathbf{N}$. Clearly, if there exists $n_0 = 1, 2, \dots$ for which $\vartheta_{n_0+1} = \vartheta_{n_0}$, then $H\vartheta_{n_0} = \vartheta_{n_0}$ and the proof is completed. Hence, we suppose that $\vartheta_{n+1} \neq \vartheta_n$ or $d_F(H\vartheta_{n-1}, H\vartheta_n) > 0$ for every $n \in \mathbf{N}$. Now as H is generalized $(\alpha\beta-\psi)$ -contraction, so we have

$$\begin{aligned} d_F(\vartheta_n, \vartheta_n) &= d_F(H\vartheta_{n-1}, H\vartheta_n) \\ &\leq \psi (\max \{d_F(\vartheta_{n-1}, \vartheta_n), \min \{d_F(\vartheta_{n-1}, H\vartheta_{n-1}), d_F(\vartheta_n, H\vartheta_n)\}\}), \end{aligned}$$

for all $n \in \mathbf{N}$. Now if $d_F(\vartheta_{n-1}, H\vartheta_{n-1}) < d_F(\vartheta_n, H\vartheta_n)$, then

$$\max \{d_F(\vartheta_{n-1}, \vartheta_n), \min \{d_F(\vartheta_{n-1}, H\vartheta_{n-1}), d_F(\vartheta_n, H\vartheta_n)\}\} = d_F(\vartheta_{n-1}, \vartheta_n),$$

for all $n \in \mathbf{N}$. If $d_F(\vartheta_n, H\vartheta_n) < d_F(\vartheta_{n-1}, H\vartheta_{n-1})$, then

$$\max \{d_F(\vartheta_{n-1}, \vartheta_n), \min \{d_F(\vartheta_{n-1}, H\vartheta_{n-1}), d_F(\vartheta_n, H\vartheta_n)\}\} = d_F(\vartheta_{n-1}, \vartheta_n),$$

for all $n \in \mathbf{N}$. Thus in all case, we have

$$d_F(\vartheta_n, \vartheta_n) \leq \psi(d_F(\vartheta_{n-1}, \vartheta_n)), \tag{4}$$

for all $n \in \mathbb{N}$. Continuing in this way, we get

$$d_F(\vartheta_n, \vartheta_n) \leq \psi^n(d_F(\vartheta_0, \vartheta_1)), \tag{5}$$

for all $n \in \mathbb{N}$. Suppose $f \in F$ and $h \in [0, +\infty)$ are such that the assertion (D₃) hold and suppose $e > 0$.

Now from (F₂), there exists $\delta > 0$ such that

$$0 < \iota < \delta \implies f(\iota) < f(\delta) - h. \tag{6}$$

Let $n(e) \in \mathbb{N}$ be such that $0 < \sum_{i \geq n(e)} \psi^i(d_F(\vartheta_0, \vartheta_1)) < \delta$. Hence, by (5), (F₁) and (F₂), we have

$$f\left(\sum_{i=n}^{m-1} d_F(\vartheta_i, \vartheta_{i+1})\right) \leq f\left(\sum_{i=n}^{m-1} \psi^i(d_F(\vartheta_0, \vartheta_1))\right) \leq f\left(\sum_{n \geq n(e)} \psi^i(d_F(\vartheta_0, \vartheta_1))\right) < f(e) - h, \tag{7}$$

for $m > n \geq n(e)$. Using (D₃) and (7), we obtain $d_F(\vartheta_n, \vartheta_m) > 0$, $m > n \geq n(e)$ implies

$$f(d_F(\vartheta_n, \vartheta_m)) \leq f\left(\sum_{i=n}^{m-1} d_F(\vartheta_i, \vartheta_{i+1})\right) + h < f(e).$$

By (F₁), we have $d_F(\vartheta_n, \vartheta_m) < e$, $m > n \geq n(e)$. This proves that $\{\vartheta_n\}$ is F-Cauchy. Since (M, d_F) is F-complete, so $\exists \vartheta^* \in M$ such that

$$\lim_{n \rightarrow \infty} d_F(\vartheta_n, \vartheta^*) = 0. \tag{8}$$

Secondly as $\vartheta_n \rightarrow \vartheta^*$ and $\alpha(\vartheta_n, \vartheta_{n+1}) \geq \beta(\vartheta_n, \vartheta_{n+1})$, then $\alpha(\vartheta^*, H\vartheta^*) \geq \beta(\vartheta^*, H\vartheta^*)$. Thus

$$\alpha(\vartheta^*, H\vartheta^*)\alpha(\vartheta_n, H\vartheta_n) \geq \beta(\vartheta^*, H\vartheta^*)\beta(\vartheta_n, H\vartheta_n).$$

We start with contradiction by supposing that $d_F(H(\vartheta^*), \vartheta^*) > 0$. By (F₁) and (D₃), we get

$$\begin{aligned} f(d_F(H(\vartheta^*), \vartheta^*)) &\leq f(d_F(H(\vartheta^*), H(\vartheta_n)) + d_F(H(\vartheta_n), \vartheta^*)) + h \\ &\leq f(d_F(H(\vartheta^*), H(\vartheta_n)) + d_F(H(\vartheta_n), \vartheta^*)) + h. \end{aligned}$$

By (1), we have

$$\begin{aligned} f(d_F(H(\vartheta^*), \vartheta^*)) &\leq f(d_F(H(\vartheta^*), H(\vartheta_n)) + d_F(H(\vartheta_n), \vartheta^*)) + h \\ &\leq f(\psi(\max\{d_F(\vartheta^*, \vartheta_n), \min(d_F(\vartheta^*, H(\vartheta^*)), d_F(\vartheta_n, H(\vartheta_n)))\} \\ &\quad + d_F(\vartheta_{n+1}, \vartheta^*))) + h, \\ &= f(\psi(\max\{D(\vartheta^*, \vartheta_n), \min(d_{\mathcal{F}}(\vartheta^*, \mathcal{H}(\vartheta^*)), d_{\mathcal{F}}(\vartheta_n, \vartheta_{n+1}))\})) \end{aligned}$$

$$+d_F(\vartheta_{n+1}, \vartheta^*) + h,$$

for all $n \in \mathbb{N}$. Letting $n \rightarrow \infty$ and using (F₂) and (8), we get

$$\lim_{n \rightarrow \infty} f(d_F(H(\vartheta^*), \vartheta^*)) \leq n \lim_{n \rightarrow \infty} f(d_F(\vartheta^*, \vartheta_n) + D(\vartheta_{n+1}, \vartheta^*) + h) = -\infty.$$

This implies that $d_F(H(\vartheta^*), \vartheta^*) = 0$, which is a contradiction.

Thus $d_F(H(\vartheta^*), \vartheta^*) = 0$, that is, $H(\vartheta^*) = \vartheta^*$. As consequence, $\vartheta^* \in \mathbf{M}$ is the fixed point of H .

□

Example 2. Let $M = \mathbb{R}$ endowed with F-metric d_F given by

$$d_F(\vartheta, \theta) = \begin{cases} e^{|\vartheta - \theta|}, & \text{if } \vartheta \neq \theta \\ 0, & \text{if } \vartheta = \theta. \end{cases}$$

Then (M, d_F) is F-complete F-metric space with $f(t) = \frac{-1}{t}$ and $h = 1$. Define $H : M \rightarrow M$ and

$$\alpha, \beta : M \times M \rightarrow [0, \infty) \text{ by } \mathcal{H}(\vartheta) = \begin{cases} \frac{\vartheta}{4} & \text{if } 0 \leq \vartheta \leq 1 \\ 3\vartheta, & \text{if } \vartheta > 1 \end{cases}$$

and

$$\alpha(\vartheta, \theta) = \begin{cases} 0, & \text{otherwise} \\ 1, & \text{if } \vartheta, \theta \in [0, 1] \end{cases}$$

$$\alpha(\vartheta, \theta) = \beta(\vartheta, \theta) =$$

$$\theta =$$

$$0, \text{ otherwise.}$$

Clearly, H is generalized $(\alpha\beta\text{-}\psi)$ -contraction mapping with $\psi(\vartheta) = \frac{\vartheta}{2}$ for all $\vartheta \geq 0$ that is

$$d_F(H\vartheta, H\theta) \leq \psi(\max\{d_F(\vartheta, \theta), \min\{d_F(\vartheta, H\vartheta), d_F(\theta, H\theta)\}\}).$$

Moreover, there exists $\vartheta_0 \in \mathbf{M}$ such that $\alpha(\vartheta_0, H\vartheta_0) = 1 = \beta(\vartheta_0, H\vartheta_0)$ and H is an α -admissible mapping with respect to β . Thus all the hypotheses of

Theorem 3 are satisfied. Consequently $H\vartheta = 0$.

Corollary 1. Let (M, d_F) be a an F-metric space and let $H : M \rightarrow M$. Assume that the following assertions hold:

- (i) (M, d_F) is F-complete,
- (ii) H is an α -admissible mapping, (iii) if for $\vartheta, \theta \in \mathbf{M}$ and $\psi \in \Psi$ such that $\alpha(\vartheta, H\vartheta)\alpha(\theta, H\theta) \geq 1 \Rightarrow d_F(H\vartheta, H\theta) \leq \psi(\max\{d_F(\vartheta, \theta), \min\{d_F(\vartheta, H\vartheta), d_F(\theta, H\theta)\}\})$,

(iv) there exists $\vartheta_0 \in \mathbf{M}$ such that $\alpha(\vartheta_0, H\vartheta_0) \geq 1$,

(v) either H is an continuous or if $\{\vartheta_n\}$ is a sequence in M such that $\vartheta_n \rightarrow \vartheta$, $\alpha(\vartheta_n, \vartheta_{n+1}) \geq$

1, then $\alpha(\vartheta, H\vartheta) \geq 1$.

Then there exists $\vartheta^* \in \mathbf{M}$ such that $\vartheta^* = H\vartheta^*$.

Proof. Consider $\beta: M \times M \rightarrow [0, +\infty)$ as $\beta(\vartheta, \theta) = 1$ for all $\vartheta, \theta \in M$ in Theorem 3. \square

The following corollaries are direct consequences of Theorem 3.

Corollary 2. Let (M, d_F) be a an F-metric space and let $H : M \rightarrow M$. Assume that the following assertions hold:

- (i) (M, d_F) is F-complete,
- (ii) H is an α -admissible mapping,
- (iii) if for $\vartheta, \theta \in M$ and $\psi \in \Psi$ such that

$$(d_F(H\vartheta, H\theta) + l)^{\alpha(\vartheta, H\vartheta)\alpha(\theta, H\theta)} \leq \psi (\max \{d_F(\vartheta, \theta), \min \{d_F(\vartheta, H\vartheta), d_F(\theta, H\theta)\}\}) + l,$$

where $l > 0$,

- (iv) there exists $\vartheta_0 \in M$ such that $\alpha(\vartheta_0, H\vartheta_0) \geq 1$,
- (v) either H is an continuous or if $\{\vartheta_n\}$ is a sequence in M such that $\vartheta_n \rightarrow \vartheta$, $\alpha(\vartheta_n, \vartheta_{n+1}) \geq$

1, then $\alpha(\vartheta, H\vartheta) \geq 1$.

Then there exists $\vartheta^* \in M$ such that $\vartheta^* = H\vartheta^*$.

Corollary 3. Let (M, d_F) be a an F-metric space and let $H : M \rightarrow M$. Assume that the following assertions hold:

- (i) (M, d_F) is F-complete,
- (ii) H is an α -admissible mapping,
- (iii) if for $\vartheta, \theta \in M$ and $\psi \in \Psi$ such that

$$(\alpha(\vartheta, H\vartheta)\alpha(\theta, H\theta) + 1)d^F(H\vartheta, H\theta) \leq 2\psi(\max \{d_F(\vartheta, \theta), \min \{d_F(\vartheta, H\vartheta), d_F(\theta, H\theta)\}\}),$$

- (iv) there exists $\vartheta_0 \in M$ such that $\alpha(\vartheta_0, H\vartheta_0) \geq 1$,
- (v) either H is an continuous or if $\{\vartheta_n\}$ is a sequence in M such that $\vartheta_n \rightarrow \vartheta$, $\alpha(\vartheta_n, \vartheta_{n+1}) \geq$

1, then $\alpha(\vartheta, H\vartheta) \geq 1$.

Then there exists $\vartheta^* \in M$ such that $\vartheta^* = H\vartheta^*$.

Corollary 4. Let (M, d_F) be a an F-metric space and let $H : M \rightarrow M$. Assume that the following assertions hold:

- (i) (M, d_F) is F-complete,
- (ii) H is an α -admissible mapping,
- (iii) if for $\vartheta, \theta \in M$ and $\psi \in \Psi$ such that $\alpha(\vartheta, H\vartheta)\alpha(\theta, H\theta)d_F(H\vartheta, H\theta) \leq k \max \{d_F(\vartheta, \theta), \min$

$$\{d_F(\vartheta, H\vartheta), d_F(\theta, H\theta)\}\},$$

- (iv) there exists $\vartheta_0 \in M$ such that $\alpha(\vartheta_0, H\vartheta_0) \geq 1$,
- (v) either H is an continuous or if $\{\vartheta_n\}$ is a sequence in M such that $\vartheta_n \rightarrow \vartheta$, $\alpha(\vartheta_n, \vartheta_{n+1}) \geq 1$,
then $\alpha(\vartheta, H\vartheta) \geq 1$.

Then there exists $g^* \in \mathbf{M}$ such that $g^* = Hg^*$.

If $\alpha(g, \theta) = 1$, then we have the following corollaries.

Corollary 5. Let (M, d_F) be a an F-metric space and let $H : M \rightarrow M$. Assume that the following assertions hold:

- (i) (M, d_F) is F-complete,
- (ii) H is an β -subadmissible mapping, (iii) if for $g, \theta \in \mathbf{M}$ and $\psi \in \Psi$ such that $\beta(g, Hg)\beta(\theta, H\theta) \leq 1 \Rightarrow d_F(Hg, H\theta) \leq \psi (\max \{d_F(g, \theta), \min \{d_F(g, Hg), d_F(\theta, H\theta)\}\})$,
- (iv) there exists $g_0 \in \mathbf{M}$ such that $\beta(g_0, Hg_0) \leq 1$,
- (v) either H is an continuous or if $\{g_n\}$ is a sequence in M such that $g_n \rightarrow g, \beta(g_n, g_{n+1}) \leq 1$, then $\beta(g, Hg) \leq 1$.

Then there exists $g^* \in \mathbf{M}$ such that $g^* = Hg^*$.

Corollary 6. Let (M, d_F) be a an F-metric space and let $H : M \rightarrow M$. Assume that the following assertions hold:

- (i) (M, d_F) is F-complete,
- (ii) H is an β -subadmissible mapping,
- (iii) if for $g, \theta \in \mathbf{M}$ and $\psi \in \Psi$ such that $d_F(Hg, H\theta) + l \leq [\psi (\max \{d_F(g, \theta), \min \{d_F(g, Hg), d_F(\theta, H\theta)\}\}) + l]^{\beta(g, Hg)\beta(\theta, H\theta)}$,

where $l > 0$,

- (iv) there exists $g_0 \in \mathbf{M}$ such that $\beta(g_0, Hg_0) \leq 1$,
- (v) either H is an continuous or if $\{g_n\}$ is a sequence in M such that $g_n \rightarrow g, \beta(g_n, g_{n+1}) \leq 1$, then $\beta(g, Hg) \leq 1$.

Then there exists $g^* \in \mathbf{M}$ such that $g^* = Hg^*$.

Corollary 7. Let (M, d_F) be a an F-metric space and let $H : M \rightarrow M$. Assume that the following assertions hold:

- (i) (M, d_F) is F-complete,
- (ii) H is an β -subadmissible mapping,
- (iii) if for $g, \theta \in \mathbf{M}$ and $\psi \in \Psi$ such that

$$2d_F(Hg, H\theta) \leq (\beta(g, Hg)\beta(\theta, H\theta) + 1)\psi(\max \{d^F(g, \theta), \min \{d^F(g, Hg), d^F(\theta, H\theta)\}\}),$$

- (iv) $\exists g_0 \in \mathbf{M}$ such that $\beta(g_0, Hg_0) \leq 1$,
- (v) either H is an continuous or if $\{g_n\}$ is a sequence in M such that $g_n \rightarrow g, \beta(g_n, g_{n+1}) \leq 1$, then $\beta(g, Hg) \leq 1$.

Then there exists $g^* \in \mathbf{M}$ such that $g^* = Hg^*$.

Corollary 8. Let (M, d_F) be a an F-metric space and let $H : M \rightarrow M$. Assume that the following assertions hold:

- (i) (M, d_F) is F-complete,
- (ii) H is an β -subadmissible mapping,
- (iii) if for $\vartheta, \theta \in M$ and $\psi \in \Psi$ such that $d_F(H\vartheta, H\theta) \leq \psi (\beta(\vartheta, H\vartheta)\beta(\theta, H\theta) \max \{d_F(\vartheta, \theta), \min \{d_F(\vartheta, H\vartheta), d_F(\theta, H\theta)\}\})$,
- (iv) there exists $\vartheta_0 \in M$ such that $\beta(\vartheta_0, H\vartheta_0) \leq 1$,
- (v) either H is an continuous or if $\{\vartheta_n\}$ is a sequence in M such that $\vartheta_n \rightarrow \vartheta$, $\beta(\vartheta_n, \vartheta_{n+1}) \leq 1$, then $\beta(\vartheta, H\vartheta) \leq 1$.

Then there exists $\vartheta^* \in M$ such that $\vartheta^* = H\vartheta^*$.

Corollary 9. Let (M, d_F) be a an F-metric space and let $H : M \rightarrow M$. Assume that the following assertions hold:

- (i) (M, d_F) is F-complete,
- (ii) H is an α -admissible mapping, (ii) if for $\vartheta, \theta \in M$ and $\psi \in \Psi$ such that

$$\alpha(\vartheta, H\vartheta)\alpha(\theta, H\theta) \geq 1 \Rightarrow d_F(H\vartheta, H\theta) \leq \psi (d_F(\vartheta, \theta)),$$

- (iv) there exists $\vartheta_0 \in M$ such that $\alpha(\vartheta_0, H\vartheta_0) \geq 1$,
- (v) either H is an continuous or if $\{\vartheta_n\}$ is a sequence in M such that $\vartheta_n \rightarrow \vartheta$, $\alpha(\vartheta_n, \vartheta_{n+1}) \geq 1$, then $\alpha(\vartheta, H\vartheta) \geq 1$.

Then there exists $\vartheta^* \in M$ such that $\vartheta^* = H\vartheta^*$.

Corollary 10. Let (M, d_F) be a an F-metric space and let $H : M \rightarrow M$. Assume that the following assertions hold:

- (i) (M, d_F) is F-complete,
- (ii) H is an α -admissible mapping,
- (ii) if for $\vartheta, \theta \in M$ and $\psi \in \Psi$ such that

$$(\alpha(\vartheta, H\vartheta)\alpha(\theta, H\theta) + 1)d_F(H\vartheta, H\theta) \leq 2\psi(d_F(\vartheta, \theta)),$$

- (iv) there exists $\vartheta_0 \in M$ such that $\alpha(\vartheta_0, H\vartheta_0) \geq 1$,
- (v) either H is an continuous or if $\{\vartheta_n\}$ is a sequence in M such that $\vartheta_n \rightarrow \vartheta$, $\alpha(\vartheta_n, \vartheta_{n+1}) \geq 1$, then $\alpha(\vartheta, H\vartheta) \geq 1$.

Then there exists $\vartheta^* \in M$ such that $\vartheta^* = H\vartheta^*$.

Corollary 11. Let (M, d_F) be a an F-metric space and let $H : M \rightarrow M$. Assume that the following assertions hold:

- (i) (M, d_F) is F-complete,
(ii) H is an α -admissible mapping, (iii) if for $\vartheta, \theta \in M$ and $\psi \in \Psi$ such that

$$\alpha(\vartheta, H\vartheta)\alpha(\theta, H\theta)d_F(H\vartheta, H\theta) \leq \psi(d_F(\vartheta, \theta)),$$

(iv) there exists $\vartheta_0 \in M$ such that $\alpha(\vartheta_0, H\vartheta_0) \geq 1$,

(v) either H is a continuous mapping or if $\{\vartheta_n\}$ is a sequence in M such that $\vartheta_n \rightarrow \vartheta$, $\alpha(\vartheta_n, \vartheta_{n+1}) \geq 1$, then $\alpha(\vartheta, H\vartheta) \geq 1$.

Then there exists $\vartheta^* \in M$ such that $\vartheta^* = H\vartheta^*$.

Corollary 12 ([3]). Let (M, d_F) be an F-metric space and let $H : M \rightarrow M$ be continuous. Assume that the following assertions hold:

- (i) (M, d_F) is F-complete, (ii) if for $\vartheta, \theta \in M$ and $\psi \in \Psi$ such that

$$d_F(H\vartheta, H\theta) \leq \psi(d_F(\vartheta, \theta)).$$

Then there exists $\vartheta^* \in M$ such that $\vartheta^* = H\vartheta^*$.

Proof. Taking $\alpha(\vartheta, \theta) = 1$, for all $\vartheta, \theta \in M$ in the Corollary 11. \square

Corollary 13. Let (M, d_F) be an F-metric space and let $H : M \rightarrow M$. Assume that the following assertions hold:

- (i) (M, d_F) is F-complete,
(ii) H is an α -admissible mapping, (iii) if there exists $k \in (0, 1)$ such that

$$\alpha(\vartheta, H\vartheta)\alpha(\theta, H\theta)d_F(H\vartheta, H\theta) \leq kd_F(\vartheta, \theta),$$

for all $\vartheta, \theta \in M$

(iv) there exists $\vartheta_0 \in M$ such that $\alpha(\vartheta_0, H\vartheta_0) \geq 1$,

(v) either H is a continuous mapping or if $\{\vartheta_n\}$ is a sequence in M such that $\vartheta_n \rightarrow \vartheta$, $\alpha(\vartheta_n, \vartheta_{n+1}) \geq 1$, then $\alpha(\vartheta, H\vartheta) \geq 1$.

Then there exists $\vartheta^* \in M$ such that $\vartheta^* = H\vartheta^*$.

Corollary 14 ([2]). Let (M, d_F) be an F-metric space and let $H : M \rightarrow M$ be a continuous mapping. Assume that the following assertions hold:

- (i) (M, d_F) is F-complete, (ii) if there exists $k \in (0, 1)$ such that

$$d_F(H\vartheta, H\theta) \leq kd_F(\vartheta, \theta),$$

for all $\vartheta, \theta \in \mathbf{M}$.

Then there exists $\vartheta^* \in \mathbf{M}$ such that $\vartheta^* = H\vartheta^*$.

Proof. Taking $\alpha(\vartheta, \theta) = 1$, for all $\vartheta, \theta \in \mathbf{M}$ in the Corollary 13. \square

3. Consequences

The following results are direct consequences of main results by taking $f(i) = \ln(i)$ and $h = \ln(1)$.

Theorem 4. Let (\mathbf{M}, d) be a complete metric space and let $H : \mathbf{M} \rightarrow \mathbf{M}$. Suppose that the following assertions hold:

(i) there exist two functions $\alpha, \beta : \mathbf{M} \times \mathbf{M} \rightarrow [0, \infty)$ and $k \in [0, 1)$ such that $\alpha(\vartheta, H\vartheta)\alpha(\theta, H\theta) \geq \beta(\vartheta, H\vartheta)\beta(\theta, H\theta)$ implies

$$d(H\vartheta, H\theta) \leq k \max \{d(\vartheta, \theta), \min \{d(\vartheta, H\vartheta), d(\theta, H\theta)\}\}$$

(ii) H is an α -admissible mapping with respect to β ,

(iii) there exists $\vartheta_0 \in \mathbf{M}$ such that $\alpha(\vartheta_0, H\vartheta_0) \geq \beta(\vartheta_0, H\vartheta_0)$,

(iv) either H is a continuous or if $\{\vartheta_n\}$ is a sequence in \mathbf{M} such that $\vartheta_n \rightarrow \vartheta$, $\alpha(\vartheta_n, \vartheta_{n+1}) \geq \beta(\vartheta_n, \vartheta_{n+1})$, then $\alpha(\vartheta, H\vartheta) \geq \beta(\vartheta, H\vartheta)$.

Then there exists $\vartheta^* \in \mathbf{M}$ such that $\vartheta^* = H\vartheta^*$.

Proof. Taking $\psi(i) = ki$, where $k \in [0, 1)$ in Theorem 3. \square

Corollary 15. Let (\mathbf{M}, d) be a complete metric space and let $H : \mathbf{M} \rightarrow \mathbf{M}$. Assume that the following assertions hold:

(i) H is an α -admissible mapping,

(ii) if for $\vartheta, \theta \in \mathbf{M}$ such that $\alpha(\vartheta, H\vartheta)\alpha(\theta, H\theta) \geq 1$ implies

$$d(H\vartheta, H\theta) \leq k \max \{d(\vartheta, \theta), \min \{d(\vartheta, H\vartheta), d(\theta, H\theta)\}\},$$

(iii) there exists $\vartheta_0 \in \mathbf{M}$ such that $\alpha(\vartheta_0, H\vartheta_0) \geq 1$,

(iv) either H is a continuous or if $\{\vartheta_n\}$ is a sequence in \mathbf{M} such that $\vartheta_n \rightarrow \vartheta$, $\alpha(\vartheta_n, \vartheta_{n+1}) \geq 1$, then $\alpha(\vartheta, H\vartheta) \geq 1$.

Then there exists $\vartheta^* \in \mathbf{M}$ such that $\vartheta^* = H\vartheta^*$.

Proof. Taking $\beta(\vartheta, \theta) = 1$ in above corollary. \square

Example 3. Let $\mathbf{M} = [0, \infty)$ be endowed with the usual metric $d(\vartheta, \theta) = |\vartheta - \theta|$ for all $\vartheta, \theta \in \mathbf{M}$ and let $H : \mathbf{M} \rightarrow \mathbf{M}$ be defined by $H\vartheta = \frac{\vartheta}{4}$. Also, define $\alpha : \mathbf{M} \times \mathbf{M} \rightarrow [0, \infty)$ by $\alpha(\vartheta, \theta) = 3$ and $\exists k = \frac{1}{2} \in (0, 1)$.

Clearly, H is an α -admissible mapping. Also, $\alpha(\vartheta, H\vartheta)\alpha(\theta, H\theta) = 9 \geq 1$ for all $\vartheta, \theta \in \mathbf{M}$. Hence

$$d(H\vartheta, H\theta) = \frac{1}{4}|\vartheta - \theta| \leq \frac{1}{2} \max \left\{ |\vartheta - \theta|, \min \left\{ \left| \vartheta - \frac{\vartheta}{4} \right|, \left| \theta - \frac{\theta}{4} \right| \right\} \right\}.$$

Then the conditions of Corollary 15 hold and H has a fixed point which is 0.

Corollary 16. Let (M, d) be a complete metric space and let $H : M \rightarrow M$. Assume that the following assertions hold:

- (i) H is an α -admissible mapping,
- (ii) if for $\vartheta, \theta \in M$ such that

$$(d(H\vartheta, H\theta) + l)^{\alpha(\vartheta, H\vartheta)\alpha(\theta, H\theta)} \leq k \max \{d(\vartheta, \theta), \min \{d(\vartheta, H\vartheta), d(\theta, H\theta)\}\} + l,$$

where $l > 0$,

- (iii) there exists $\vartheta_0 \in M$ such that $\alpha(\vartheta_0, H\vartheta_0) \geq 1$,
- (iv) either H is a continuous mapping or if $\{\vartheta_n\}$ is a sequence in M such that $\vartheta_n \rightarrow \vartheta$, $\alpha(\vartheta_n, \vartheta_{n+1}) \geq 1$, then $\alpha(\vartheta, H\vartheta) \geq 1$.

Then there exists $\vartheta^* \in M$ such that $\vartheta^* = H\vartheta^*$.

Corollary 17. Let (M, d) be a complete metric space and let $H : M \rightarrow M$. Assume that the following assertions hold:

- (i) H is an α -admissible mapping,
- (ii) if for $\vartheta, \theta \in M$ such that

$$(\alpha(\vartheta, H\vartheta)\alpha(\theta, H\theta) + 1)d(H\vartheta, H\theta) \leq 2k \max \{d(\vartheta, \theta), \min \{d(\vartheta, H\vartheta), d(\theta, H\theta)\}\},$$

- (iii) there exists $\vartheta_0 \in M$ such that $\alpha(\vartheta_0, H\vartheta_0) \geq 1$,
- (iv) either H is a continuous mapping or if $\{\vartheta_n\}$ is a sequence in M such that $\vartheta_n \rightarrow \vartheta$, $\alpha(\vartheta_n, \vartheta_{n+1}) \geq 1$, then $\alpha(\vartheta, H\vartheta) \geq 1$.

Then there exists $\vartheta^* \in M$ such that $\vartheta^* = H\vartheta^*$.

Corollary 18. Let (M, d) be a complete metric space and let $H : M \rightarrow M$. Assume that the following assertions hold:

- (i) H is an α -admissible mapping,
- (ii) if for $\vartheta, \theta \in M$ such that $\alpha(\vartheta, H\vartheta)\alpha(\theta, H\theta)d(H\vartheta, H\theta) \leq k \max \{d(\vartheta, \theta), \min \{d(\vartheta, H\vartheta), d(\theta, H\theta)\}\}$,

- (iii) there exists $\vartheta_0 \in M$ such that $\alpha(\vartheta_0, H\vartheta_0) \geq 1$,
- (iv) either H is a continuous mapping or if $\{\vartheta_n\}$ is a sequence in M such that $\vartheta_n \rightarrow \vartheta$, $\alpha(\vartheta_n, \vartheta_{n+1}) \geq 1$, then $\alpha(\vartheta, H\vartheta) \geq 1$.

Then there exists $\vartheta^* \in M$ such that $\vartheta^* = H\vartheta^*$.

If $\alpha(\vartheta, \theta) = 1$, then we have the following corollaries.

Corollary 19. Let (M, d) be a complete metric space and let $H : M \rightarrow M$. Assume that the following assertions hold:

- (i) H is an β -subadmissible mapping,
- (ii) if for $\vartheta, \theta \in M$ such that $\beta(\vartheta, H\vartheta)\beta(\theta, H\theta) \leq 1$ implies

$$d(H\vartheta, H\theta) \leq k \max \{d(\vartheta, \theta), \min \{d(\vartheta, H\vartheta), d(\theta, H\theta)\}\},$$

- (iii) there exists $\vartheta_0 \in M$ such that $\beta(\vartheta_0, H\vartheta_0) \leq 1$,
- (iv) either H is an continuous or if $\{\vartheta_n\}$ is a sequence in M such that $\vartheta_n \rightarrow \vartheta, \beta(\vartheta_n, \vartheta_{n+1}) \leq 1$, then $\beta(\vartheta, H\vartheta) \leq 1$.

Then there exists $\vartheta^* \in M$ such that $\vartheta^* = H\vartheta^*$.

Corollary 20. Let (M, d) be a complete metric space and let $H : M \rightarrow M$. Assume that the following assertions hold:

- (i) H is an β -subadmissible mapping,
- (ii) if for $\vartheta, \theta \in M$ such that $d(H\vartheta, H\theta) + l \leq [k \max \{d(\vartheta, \theta), \min \{d(\vartheta, H\vartheta), d(\theta, H\theta)\}\} +$

$$\beta^{\beta(\vartheta, H\vartheta)\beta(\theta, H\theta)},$$

where $l > 0$.

- (iii) there exists $\vartheta_0 \in M$ such that $\beta(\vartheta_0, H\vartheta_0) \leq 1$,
- (iv) either H is an continuous or if $\{\vartheta_n\}$ is a sequence in M such that $\vartheta_n \rightarrow \vartheta, \beta(\vartheta_n, \vartheta_{n+1}) \leq 1$, then $\beta(\vartheta, H\vartheta) \leq 1$.

Then there exists $\vartheta^* \in M$ such that $\vartheta^* = H\vartheta^*$.

Remark 2. One can easily derive the main results of References [11] and [15] from our Corollaries 11 and 14 respectively by taking $f(\cdot) = \ln(\cdot)$ and $h = \ln(1)$.

4. Applications

In the present section, we solve the following differential equation

$$\vartheta'(t) = -\rho_1(t)\vartheta(t) + \rho_2(t)L(\vartheta(t - \zeta(t))) + \rho_3(t)\vartheta'(t - \zeta(t)). \tag{9}$$

The lemma of Djoudi et al. [16] is very handy in the proof our theorem.

Lemma 2 ([16]). Suppose that $\zeta'(t) \neq 1 \forall t \in \mathbb{R}$. Then $\vartheta(t)$ is a solution of (9) iff

$$\vartheta(t) = \left(\vartheta(0) - \frac{\rho_3(0)}{1 - \zeta'(0)} \vartheta(-\zeta(0)) \right) \rho^{-\int_0^t \alpha(\zeta) d\zeta} + \frac{\rho_3(t)}{1 - \zeta'(t)} \vartheta(t - \zeta(t))$$

$$- \int_0^t (h(\omega)\vartheta(\omega - \zeta(\omega))) - \rho_2(\omega)L(\vartheta(\omega - \zeta(\omega))) \rho^{-\int_0^t \alpha(\zeta) d\zeta} d\omega, \tag{10}$$

where

$$h(\omega) = \frac{\zeta''(\omega)\rho_3(\omega) + (\rho_3'(\omega) + \rho_3(\omega)\rho_1(\omega))(1 - \zeta'(\omega))}{(1 - \zeta'(\omega))^2}. \tag{11}$$

Now assume that $\varphi : (-\infty, 0] \rightarrow \mathbb{R}$ is a continuous bounded initial function, then $g(t) = g(t, 0, \varphi)$ is a solution of (9) if $g(t) = \varphi(t)$ for $t \leq 0$ and assures (9) for $t \geq 0$. Assume C be the collection of $\pi : \mathbb{R} \rightarrow \mathbb{R}$ which are continuous functions. Define B_φ by

$$B_\varphi = \{ \pi : \mathbb{R} \rightarrow \mathbb{R} \text{ such that } \varphi(t) = \pi(t) \text{ if } t \leq 0, \pi(t) \rightarrow 0 \text{ as } t \rightarrow \infty, \pi \in C \} .$$

Then B_φ is a Banach space equipped with the supremum norm $k \cdot k$.

Lemma 3 ([3]). The space $(B_\varphi, k \cdot k)$ provided with d given by

$$d(t, i^*) = \|t - i^*\| = \sup_{\vartheta \in I} |t(\vartheta) - i^*(\vartheta)|$$

for $t, i^* \in B_\varphi$, is an F-metric space.

Theorem 5. Let $H : B_\varphi \rightarrow B_\varphi$ be a mapping defined by

$$(H\pi)(t) = \left(\pi(0) - \frac{\rho_3(0)}{1 - \zeta'(0)} \pi(-\zeta(0)) \right) \rho^{-\int_0^t \alpha(\zeta) d\zeta} + \frac{\rho_3(t)}{1 - \zeta'(t)} \pi(t - \zeta(t))$$

$$- \int_0^t (h(\omega)\pi(\omega - \zeta(\omega)) - \rho_2(\omega)L(\pi(\omega - \zeta(\omega)))) \rho^{-\int_\omega^t \alpha(\zeta) d\zeta} d\omega, t \geq 0 \tag{12}$$

for all $\pi \in B_\varphi$. Assume that these assertions are satisfied:

(i) there exists $\mu \geq 0$ and $\varphi \in \Psi$ so that

$$\begin{aligned} & \int_0^t |h(\omega)(\pi(\omega - \zeta(\omega))) - \mathcal{S}(\omega - \zeta(\omega))| \rho^{-\int_\omega^t \alpha(\zeta) d\zeta} \\ & \leq -\mu_\varphi (\max \{k_\pi - \mathcal{S}k, \min \{k_\pi - H\pi k, k\mathcal{S} - H\mathcal{S}k\}\}) \end{aligned} \tag{13}$$

and

$$\begin{aligned} & \int_0^t |(\rho_2(\omega))\mathcal{L}(\pi(\omega - \zeta(\omega))) - \mathcal{L}(\varrho(\omega - \zeta(\omega)))| \rho^{-\int_\omega^t \alpha(\zeta) d\zeta} \\ & \leq -\mu_\varphi (\max \{k_\pi - \mathcal{S}k, \min \{k_\pi - H_\pi k, k_\mathcal{S} - H_\mathcal{S}k\}\}) \end{aligned} \tag{14}$$

for all $\pi, \mathcal{S} \in B_\varphi$.

(ii) Then H has a fixed point.

Proof. Define $\alpha : C \times C \rightarrow \mathbb{R}$ by

$$\alpha(\pi, \mathcal{S}) = \beta(\pi, \mathcal{S}) = \begin{cases} 1, & \text{if } \pi, \mathcal{S} \in B_\varphi, \\ 0, & \text{otherwise.} \end{cases}$$

Now for $\pi, \mathcal{S} \in B_\varphi$ such that $\alpha(\pi, \mathcal{S}) = \beta(\pi, \mathcal{S}) \geq 1$. It follows from (12) that $H(\pi), H(\mathcal{S}) \in B_\varphi$.

Therefore $\alpha(H(\pi), H(\mathcal{S})) = \beta(H(\pi), H(\mathcal{S})) \geq 1$. Since, (13)–(15) hold, then for $\pi, \mathcal{S} \in B_\varphi$, we have

$$\begin{aligned}
|(\mathcal{H}\pi)(t) - (\mathcal{H}q)(t)| &\leq \left| \frac{\rho_3(t)}{1 - \zeta'(t)} \right| \|\pi - q\| \\
&\quad + \int_0^t |h(\omega)(\pi(\omega - \zeta(\omega))) - q(\omega - \zeta(\omega))| \rho^{-\int_0^t \alpha(\zeta) d\zeta} \\
&\quad \int_0^t |(\rho_2(\omega))\mathcal{L}(\pi(\omega - \zeta(\omega))) - \mathcal{L}(q(\omega - \zeta(\omega)))| \rho^{-\int_0^t \alpha(\zeta) d\zeta} \\
&\leq \left| \frac{\rho_3(t)}{1 - \zeta'(t)} \right| \|\pi - q\| + \mu \phi(\max\{\|\pi - q\|, \min\{\|\pi - \mathcal{H}\pi\|, \|q - \mathcal{H}q\|\}\}) \\
&\leq \left\{ \left| \frac{\rho_3(t)}{1 - \zeta'(t)} \right| + \mu \right\} \phi(\max\{\|\pi - q\|, \min\{\|\pi - \mathcal{H}\pi\|, \|q - \mathcal{H}q\|\}\}) \\
&\leq \phi(\max\{\|\pi - q\|, \min\{\|\pi - \mathcal{H}\pi\|, \|q - \mathcal{H}q\|\}\}).
\end{aligned}$$

Hence, $dF(\mathcal{H}\pi, \mathcal{H}q) \leq \psi(\max\{dF(\pi, q), \min\{dF(\pi, \mathcal{H}\pi), dF(q, \mathcal{H}q)\}\})$

implies that \mathcal{H} is generalized $(\alpha\beta\text{-}\psi)$ -contraction. Thus by Theorem 3, \mathcal{H} has a unique fixed point in B_ϕ which solves (9). \square

5. Conclusions

In this paper, we defined generalized $(\alpha\beta\text{-}\psi)$ -contraction in the setting of F-metric space and obtained some new fixed point results. As consequence of main results, we derived some fixed point results in metric spaces. We investigated the existence of solution for the following nonlinear neutral differential equation with an unbounded delay as application of our main results.

References

1. Frechet, M. Sur quelques points du calcul fonctionnel. *Rendiconti del Circolo Matematico di Palermo* **1906**, 22, 1–72.
2. Jleli, M.; Samet, B. On a new generalization of Metric Spaces. *J. Fixed Point Theory Appl.* **2018**, 2018, 128.
3. Hussain, A.; Kanwal, T. Existence and uniqueness for a neutral differential problem with unbounded delay via fixed point results. *Trans. A Razmadze Math. Inst.* **2018**, 172, 481–490.
4. Alnaser, L.A.; Lateef, D.; Fouad, H.A.; Ahmad, J. Relation theoretic contraction results in F-metric spaces. *J. Nonlinear Sci. Appl.* **2019**, 12, 337–344.
5. Alnaser, L.A.; Lateef, D.; Fouad, H.A.; Ahmad, J. New fixed point theorems with applications to non-linear neutral differential equations. *Symmetry* **2019**, 11, 602.
6. Lateef, D.; Ahmad, J. Dass and Gupta's Fixed point theorem in F-metric spaces. *J. Nonlinear Sci. Appl.* **2019**, 12, 405–411.
7. Al-Mazrooei, A.E.; Ahmad, J. Fixed Point Theorems for Rational Contractions in F-Metric Spaces. *J. Math. Anal.* **2019**, 10, 79–86.
8. Aydi, H.; Karapinar, E.; Mitrovic', Z.D.; Rashid, T. A remark on "Existence and uniqueness for a neutral differential problem with unbounded delay via fixed point results F-metric spaces". *Revista de la Real Academia de Ciencias Exactas Físicas y Naturales. Serie A. Matemáticas* **2019**, 113, 3197–3206.

9. Alqahtani, O.; Karapınar, E.; Shahi, P. Common fixed point results in function weighted metric spaces. *J. Inequalities Appl.* **2019**, *2019*, 164.
10. Al-Mezel, S.A.; Ahmad, J. Generalized fixed point results for almost (α, F_ϕ) -contractions with applications to Fredholm integral inclusions. *Symmetry* **2019**, *11*, 1068.
11. Samet, B.; Vetro, C.; Vetro, P. Fixed point theorems for α - ψ -contractive type mappings. *Nonlinear Anal.* **2012**, *75*, 2154–2165.
12. Hussain, N.; Salimi, P. Suzuki-Wardowski type fixed point theorems for α -GF-contractions. *Taiwan J. Math.* **2014**, *2014*, 18.
13. Ahmad, J.; Al-Rawashdeh, J.; Azam, A. New Fixed Point Theorems for Generalized F -Contractions in Complete Metric Spaces. *Fixed Point Theory Appl.* **2015**, *2015*, 80.
14. Hussain, N.; Ahmad, J.; Azam, A. On Suzuki-Wardowski type fixed point theorems. *J. Nonlinear Sci. Appl.* **2015**, *8*, 1095–1111.
15. Banach, S. Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales. *Fund. Math.* **1922**, *3*, 133–181.
16. Djoudi, A.; Khemis, R. Fixed point techniques and stability for natural nonlinear differential equations with unbounded delays. *Georgian Math. J.* **2006**, *13*, 25–34.